



- b) The project activity does not lead to displacement of production of goods or delivery of utilities;
- c) Environmental conditions and human-caused degradation do not permit the encroachment of natural forest vegetation;
- d) Lands will be afforested/reforested by adding NanoClay and direct planting or seeding, with trees/shrubs complying with the forest definition of DNA after forest establishment;
- e) Inter-cropping between rows of trees/shrubs is allowed in initial years of the project activity and will then be included in the monitoring;
- f) Nitrogen-fixing species are allowed to be used;
- g) Plantation may be harvested with either short or long rotation and will be regenerated either by direct planting or natural sprouting;
- h) Carbon stocks in litter and deadwood can be expected to decrease more or increase less in the absence of the project activity, relative to the project scenario;
- i) Grazing will not occur within the project boundary in both the project case and baseline scenario;
- j) Site preparation and intercropping may cause a significant long-term net emission from soil carbon;
- k) If the proposed A/R CDM project activity produces forage to feed livestock, all forage shall have a similar nutritional value and digestibility, and will support only a single livestock group with a single manure management system:
- l) Biomass burning for site preparation is not practiced.

**3. Selected carbon pools**

*Table 1: Selection and justification of carbon pools*

Carbon Pools	Selected (answer with yes or no)	Justification / Explanation
Above ground	Yes	Major carbon pool subjected to the project activity
Below ground	Yes	Major carbon pool subjected to the project activity
Dead wood	No	Conservative approach under applicability condition
Litter	No	Conservative approach under applicability condition
Soil organic carbon	Yes	Major carbon pool subjected to the project activity

**4. Summary of baseline and monitoring methodologies**

The methodology is applicable to AR CDM project activities on degraded and degrading land, which is either abandoned barren land or grassland. The major baseline and monitoring methodological steps are summarized below respectively.

Baseline methodology steps:

The eligibility of A/R activity as A/R CDM project activity is demonstrated according to Decision 5/CMP.1 (“Modalities and procedures for afforestation and reforestation project activities under the clean development mechanism in the first commitment period of the Kyoto Protocol”). This methodology applies approach 22(a) as a general baseline approach for the proposed A/R CDM

---

because there are prohibitive barriers to the policy in this area, etc ). If the policies (implemented before 11 Nov 2001) significantly impact the project area, then the baseline scenario cannot be “arid and hyper arid deserts” and this methodology cannot be used”.

project activity, taking into account historic land use/cover changes, national, local and sectoral policies that influence land use within the boundary of the proposed A/R CDM project activity, economical attractiveness of the project relative to the baseline, and barriers for implementing project activities in absence of CDM finance.

The proposed A/R CDM project area is stratified based on local site classification map/table, the current land use/cover maps or satellite image, soil map, vegetation map, landform map as well as supplementary surveys when necessary, and the baseline scenario is determined separately for each stratum. For strata without growing trees, this methodology conservatively assumes that the carbon stock in above-ground and below-ground biomass and soil organic matter would remain constant in the absence of the project activity, i.e., the baseline net GHG removals by sinks are zero. For strata with a few growing trees, the carbon stock change in above-ground and below-ground biomass are estimated based on methods included in GPG-LULUCF<sup>3</sup> and the carbon stock change in soil organic matter is assumed to be zero. Only the carbon stock changes in above-ground and below-ground biomass (in living trees) and soil organic matter are estimated. The omission of the dead wood and litter is considered to be conservative because it can be justified that these pools would decrease more or increase less in the absence of the proposed A/R CDM project activity, relative to the project scenario. The loss of non-tree living biomass on the site due to competition from planted trees or site preparation is accounted as an emission within the project boundary, in a conservative manner.

This methodology uses the latest version of the “Tool for the demonstration and assessment of additionality for afforestation and reforestation CDM project activities” approved by the CDM Executive Board<sup>4</sup>.

#### Monitoring methodology steps:

The monitoring methodology aims to supervise the overall project performance, including the integrity of project boundaries and forestation and management success, the actual net GHG removals by sinks, increase in GHG emissions within the project boundary due to nitrogen fertilization, planting or intercropping of N-fixing species, machinery use in site preparation, thinning and logging, and removing of existing non-tree vegetation due to site preparation or competition from planted species. It accounts for leakage due to increased domestic livestock and vehicle use for transportation staff, seedlings, timber and non forest products. A Quality Assurance/Quality Control plan, including field measurements, data collection verification, data entry and archiving, is integrated in the monitoring plan.

The baseline net GHG removals by sinks do not need to be measured and monitored over time. However, the methodology checks and re-assesses these assumptions if a renewal of the crediting period is chosen. This methodology stratifies the project area based on local climate, existing vegetation, site class and species and/or years to be planted with the aid of land use/cover maps, satellite images or aerial photograph, soil map, and field survey. This methodology uses permanent sample plots to monitor carbon stock changes in living biomass and soil organic matter pools. The methodology first determines the number of plots needed in each stratum/sub-stratum to reach the targeted precision level of  $\pm 10\%$  of the mean at the 95% confidence level. GPS is used to locate plots.

---

<sup>3</sup> Throughout this document, “GPG-LULUCF” means the Good Practice Guidance for Land Use, Land Use Change and Forestry from the Intergovernmental Panel on Climate Change (2003). This document is available at the following URL: <http://www.ipcc-nggip.iges.or.jp/public/gpplulucf/gpplulucf.htm>.

<sup>4</sup> Throughout this document, “A/R additionality tool” refers to the document approved by the Executive Board of the CDM and available at the following URL: <http://cdm.unfccc.int/EB/Meetings/021/eb21repan16.pdf>.

## Section II. Baseline methodology description

### 1. Eligibility of land

Eligibility of the project activities as the A/R CDM project activities under Article 12 of the Kyoto Protocol shall be demonstrated based on definitions provided in paragraph 1 of the annex to the Decision 16/CMP.1 (“Land use, land-use change and forestry”), as requested by Decision 5/CMP.1 (“Modalities and procedures for afforestation and reforestation project activities under the clean development mechanism in the first commitment period of the Kyoto Protocol”), until procedures to demonstrate the eligibility of lands for afforestation and reforestation project activities under the clean development mechanism are recommended by the EB.

### 2. Project boundary

#### *Physical delineation*

The A/R CDM project activity may contain more than one discrete parcel of land. Each discrete parcel of land shall have a unique geographical identification. The boundary shall be defined for each discrete parcel. The discrete parcels of lands will be defined by polygons, and to make the boundary geographically verifiable and transparent, information for the physical delineation of the project boundary that shall be provided, including:

- The name of the project area including name of villages, towns/townships, compartment number, allotment number, etc.
- Maps of the area showing the project boundary (paper format or digital format).
- Geographical coordinates, for each corner of the polygon sites collected using GPS, analysis of geo-referenced spatial data, or other appropriate techniques..

#### *Identification of all GHG emission sources in the project boundary*

Furthermore, the project boundary includes the emission sources and gases listed in Table 2 below.

**Table 2: GHG emissions from sources other than those resulting from changes in carbon pools within the project boundary.**

Source	Gas	Included/ excluded	Justification / Explanation
Combustion of fossil fuels	CO <sub>2</sub>	Included	Potential significant emission source
	CH <sub>4</sub>	Excluded	Potential emission is negligibly small
	N <sub>2</sub> O	Excluded	Potential emission is negligibly small
N-fixing species	CO <sub>2</sub>	Excluded	Not applicable
	CH <sub>4</sub>	Excluded	Not applicable
	N <sub>2</sub> O	Included	Potential significant emission source
Use of fertilizers	CO <sub>2</sub>	Excluded	Not applicable
	CH <sub>4</sub>	Excluded	Not applicable
	N <sub>2</sub> O	Included	Potential significant emission source
Livestock fed with forage produced by the project	CO <sub>2</sub>	Excluded	Not applicable
	CH <sub>4</sub>	Included	Potential significant emission source
	N <sub>2</sub> O	Included	Potential significant emission source

### 3. Ex-ante stratification

The stratification for ex-ante estimation of net anthropogenic GHG removals by sinks includes stratification for baseline scenario and stratification for project scenario. They can be implemented using the following steps:

#### a) Baseline stratification

**Step 1:** Assessing the key factors influencing carbon stocks in the above- and below-biomass pools. These factors may include soil features, micro-climate, landform (e.g. elevation, slope gradient), tree species to be planted, year to be planted, human management, etc.

**Step 2:** Collecting local information/information media of key factors identified in step 1, e.g.:

- local site classification maps or tables;
- the most updated land use/cover maps or satellite images / aerial photography;
- geographical, geological and soil maps visualising landform, parent rocks, soil types, and soil erosion intensity;
- other information relevant to key factors identified above.

Data sources may include archives, critically reviewed records, statistics, study reports and publications of national, regional or local governments, institutes, agencies or other authorities, and literature. Use of remote sensing products is recommended (e.g. aerial photos, satellite images, etc.).

**Step 3:** Preliminary stratification: Preferably, the stratification shall be carried out on GIS platforms by overlaying information/maps collected. If the GIS platform is not used then the stratification shall be conducted in a hierarchical order that depends on the significance of key factors on carbon stock changes or the extent of difference of the key factors across the project area. Only once higher level stratification is complete shall stratification at the next level down commence. At each level in the hierarchy, stratification shall be conducted within the strata determined at the upper level. For example, if there is a significant climatic difference within the project boundary, the stratification process may begin with stratification according to difference of the climate. If the key factor in the second level is soil type, then strata determined in the first level may be further stratified based on difference of soil type.

**Step 4:** Carrying out a supplementary sampling survey on site specifications for each preliminary stratum, e.g.:

- Existing trees if any: species, age class, number of trees, mean diameter at breast height (DBH) or height by measuring randomly selected plots with an area of 400 m<sup>2</sup> (at least three plots for each preliminary stratum);
- Non-tree vegetation: crown cover and mean height for herbaceous vegetation and shrubs by measuring randomly selected plots with an area of 1-4 m<sup>2</sup> (at least 10 plots for each preliminary stratum). For stratum with growing trees, the plots can be sub-plots of plots for measuring trees;
- Site and soil factors: soil type, soil depth, slope gradient, intensity of soil erosion, underground water level, etc. and sampling soils for soil organic matter determination;
- Human intervention: prescribed burning, logging (if any), grazing, fuel collecting, medicine collection and others;
- Conducting variation analysis for key factors investigated above. If the variation is large within each preliminary stratum, more intense field investigation shall be conducted or further stratification shall be considered in step 5.

**Step 5:** Final baseline stratification based on supplementary information collected from step 4 above, by checking whether or not each preliminary stratum is sufficiently homogenous or the difference among preliminary strata is significant. The degree of homogeneity may vary from project to project. A stratum within which there is a significant variation in any factor like vegetation type, soil type and human intervention shall be divided into two or more strata. Strata with similar features shall be merged into one stratum. Distinct strata should differ significantly from each other in terms of their numerical values for the input variables regarding baseline or project carbon calculation. For

example, sites with different species and age classes of trees already growing shall form separate strata, if, e.g. growth differences or biomass levels of species or age classes justify this. Sites with a more intensive collection of fuelwood might also be a separate stratum. However, site and soil factors may not warrant a separate stratum as long as all lands have a baseline of continued degradation, with little to no vegetation growing, and with no human intervention, and as long as the carbon accumulation in above-ground and below-ground biomass and soil organic carbon is similar in the project scenario.

**Step 6:** Visualize the stratification in a map, preferably using a Geographical Information System (GIS).

### **b) Project stratification**

Tree growth under the project scenario in the baseline strata may differ even for the same tree species at a same age class, depending on site conditions of the baseline strata. In contrast, planted trees in two or more baseline strata may have similar growth rates under the project scenario. For ex ante stratification it is preferable to conduct a project stratification based on baseline strata by including substrata, using the following steps:

**Step 1:** Define the “stand models” to be implemented in the project area by specifying:

- The species or species combination to be planted together in one single location and at the same date to create a “stand model”;
- The growth assumptions for each species or combination of species in the stand models;
- Planting, fertilization, thinning, harvesting, coppicing, and replanting cycle scheduled for each stand model, by specifying:
  - The age class when the above management activities will be implemented;
  - The quantities and types of fertilizers to be applied;
  - The volumes to be thinned or harvested;
  - The volumes to be left on site (harvest residues becoming dead wood) or extracted.

**Step 2:** Define the establishment timing of each stand model by specifying:

- The planting date;
- The area to be planted (ha);
- The geographical location for each stand model.

**Step 3:** Create sub-strata for each baseline stratum based on information collected in step 1 and step 2 above.

**Step 4:** Create a stratification map, preferably using a Geographical Information System (GIS). The GIS will be useful for integrating the data from different sources, which can then be used to identify and stratify the project area, facilitating consistency with the project boundary, and transparent monitoring and ex-post stratification. The boundary of each stratum can be delineated using GPS, analysis of geo-referenced spatial data, including remotely sensed images, or other appropriate techniques. Check the consistency with the overall project boundary.

**Notes:** In the equations used in this methodology, the letter "i" is used to represent a stratum, the letter "j" represent a species in baseline scenario and an age class (sub-stratum) in project scenario, and "k" represent species in project scenario. Ex-post adjustments of the strata in the project scenario (ex-post stratification) may be needed if unexpected disturbances occur during the crediting period (e.g. due to fire, storm, pests or disease outbreaks), affecting different parts of an originally homogeneous stratum or stand, or when forest management (planting, thinning, harvesting, replanting) occurs at different intensities, dates and spatial locations than originally planned, or if monitoring finds similar carbon stock change in different strata or significant variation in carbon stock change within one stratum / substratum.

#### 4. Procedure for selection of most plausible baseline scenario

Project participants should determine the most plausible baseline scenario with the following steps:

**Step 1:** Define the project boundary as described in Section II.2 above.

**Step 2:** Analyze historical land use, local and sectoral land-use policies or regulations

- a) Analyze the historical and existing land use/cover changes in a social-economic context and identify key factors that influence the land use/cover changes over time, using multiple sources of data including archives, maps or satellite images of land use/cover data attributable to 31.12.1989 (reforestation) or at least 50 years old (afforestation) and before the start of the proposed AR CDM project activity, supplementary field investigation, land-owner interviews, as well as studies and data collected from other sources.
- b) Demonstrate that historical and current land-use / land-cover change has lead to progressive degradation of the land over time including a decrease or steady state of the carbon stocks in the carbon pools and that it is likely that no natural encroachment of trees will occur. The demonstration should be based on verifiable information, such as scientific literature and studies or data collected in the project area or similar areas. The historical degradation feature can be indicated by assessing one of the following indicators:
  - Vegetation degradation, e.g.,
    - The crown cover of non-tree vegetation has decreased in the recent past for reasons other than sustainable harvesting activities;
  - Soil degradation, e.g.,
    - Soil erosion has increased between two time points in the recent past;
    - Soil organic matter content has decreased between two time points in the recent past.

The fact that no natural encroachment of trees would occur can be demonstrated by,

- demonstration of lack of on-site seed pool that may results in natural regeneration;
  - demonstration of lack of external seed sources that may result in natural regeneration;
  - demonstration of lack of possibility of seed sprouting and growth of young trees;
  - demonstration of lack of possible natural regeneration activity, by use of supplementary surveys on the project areas as well as similar surrounding areas for two different years that cover a minimum time period of ten years;
  - any other evidence that demonstrates the impossibility of natural encroachment in a credible and verifiable way.
- c) Demonstrate that national or sectoral land-use policies or regulations that create policy driven market distortions which give comparative advantages to afforestation/reforestation activities and that have been adopted before 11 November 2001 do not influence the areas of the proposed A/R CDM project activity (e.g., because the policy is not implemented, the policy does not target this area, or because there are prohibitive barriers to the policy in this area, etc<sup>5</sup>). If the policies (implemented before 11 Nov 2001) significantly impact the project area, then the baseline scenario cannot be “arid and hyper arid deserts” and this methodology cannot be used.

**Step 3:** Stratify the project area as explained in Section II.3 above.

**Step 4:** Identify and list plausible alternative land uses including alternative future public or private activities on the arid and hyper arid deserts such as any similar A/R activity or any other feasible land

---

<sup>5</sup> To comply with ruling of the Executive Board of the CDM, see <http://cdm.unfccc.int/EB/Meetings/016/eb16repan3.pdf>

development activities, considering relevant national or sectoral land-use policies that would impact the proposed project area, and land records, field surveys, data and feedback from stakeholders, and other appropriate sources.

**Step 5:** Demonstrate that under the plausible scenarios identified in Step 3, the most plausible scenario is that the project areas would remain degraded and degrading in absence of the project activity, by assessing the attractiveness of the plausible alternative land uses in terms of benefits to the project participants, consulting with stakeholders for existing and future land use, and identifying barriers for alternative land uses. This can be done in at least one of the following ways:

- **Generally:** By demonstrating that similar lands, in the vicinity, are also not, and are not planned to be used for these alternative land uses. Show that apparent financial or other barriers, which prevent alternative land uses can be identified;
- **Specifically for a forest as alternative land use:** Apply step 2 (investment analysis) or step 3 (barrier analysis) of the A/R “Tool for the demonstration and assessment of additionality”, to demonstrate that this land use, in absence of the CDM, is unattractive;
- **Specifically for any agricultural alternative land uses:** Demonstrate that the project lands are legally restricted to forestry purposes only, and that these restrictions are generally complied with in the vicinity of the project area, and then use the second bullet above to demonstrate that forestry land use, in absence of the CDM, is unattractive. Alternatively, use step 2 of the A/R “Tool for the demonstration and assessment of additionality” to demonstrate that alternative agricultural land uses are financially non-viable.

This methodology is not applicable if project proponents can not clearly show in the application of Steps 1 to 5 that the baseline approach 22(a) (existing or historical changes in carbon stocks in the carbon pools within the project boundary) and the scenario “lands to be planted are arid and hyper arid deserts and will continue to degrade in absence of the project” is the most appropriate plausible baseline scenario.

To ensure transparency regarding the condition of arid and hyper arid deserts, all information used in the analysis and demonstration shall be archived and verifiable.

## 5. Estimation of baseline net GHG removals by sinks

The baseline net GHG removals by sinks is the sum of the baseline net GHG removals by sinks across all strata. For those strata without trees, the sum of carbon stock changes in above-ground and below-ground biomass is set as zero. For those strata with growing trees, the sum of carbon stock changes in above-ground and below-ground biomass is determined based on the projection of their number and growth, based on growth models (yield tables), allometric equations, and local or national or IPCC default parameters (see detail below in this section). The carbon stock changes in soil organic matter for all strata in baseline scenario are set as zero. The following formulae are used to calculate the baseline net GHG removals by sinks:

$$C_{BSL,t} = \sum_i \sum_j \Delta C_{ij,BSL,t} \quad (\text{B.1})$$

where:

$i$	strata
$j$	tree species
$C_{BSL,t}$	sum of the changes in carbon stocks in trees for year t, tonnes CO <sub>2</sub> yr <sup>-1</sup>
$\Delta C_{ij,BSL,t}$	average annual carbon stock change for stratum i, species j in the absence of the project activity for year t, tonnes CO <sub>2</sub> yr <sup>-1</sup>
$t$	year 1 to length of crediting period



Carbon stock change in soil organic matter for all strata is set as zero in the baseline scenario. For those strata without growing trees,  $\Delta C_{ij,BSL,t} = 0$ . For those strata with a few growing trees,  $\Delta C_{ij,BSL,t}$  is estimated using one of following two methods for biomass growth of living trees that can be chosen based on the availability of data.

**(a) Method 1 (Carbon gain-loss method)<sup>6</sup>**

$$\Delta C_{ij,t} = (\Delta C_{G,ij,t} - \Delta C_{L,ij,t}) \quad (\text{B.2})$$

where:

$\Delta C_{ij,t}$	average annual carbon stock change due to biomass growth of living trees for stratum i, species j, tonnes CO <sub>2</sub> yr <sup>-1</sup> for year t
$\Delta C_{G,ij,t}$	average annual increase in carbon due to biomass growth of living trees for stratum i, species j, tonnes CO <sub>2</sub> yr <sup>-1</sup> for year t
$\Delta C_{L,ij,t}$	average annual decrease in carbon due to biomass loss of living trees for stratum i, species j, tonnes CO <sub>2</sub> yr <sup>-1</sup> for year t. To be conservative for the baseline scenario, $\Delta C_{L,ij} = 0$ in this methodology.

$$\Delta C_{G,ij,t} = A_i \cdot G_{TOTAL,ij,t} \cdot CF_j \cdot 44/12 \quad (\text{B.3})$$

where:

$\Delta C_{G,ij,t}$	average annual increase in carbon due to biomass growth of living trees for stratum i, species j, tonnes CO <sub>2</sub> yr <sup>-1</sup> for year t
$A_{ij}$	area of stratum i, species j, hectare (ha)
$G_{TOTAL,ij,t}$	average annual increment of total dry biomass of living trees for stratum i, species j, tonnes of dry matter, ha <sup>-1</sup> yr <sup>-1</sup> for year t
$CF_j$	the carbon fraction for species j, tonnes C (tonne d.m.) <sup>-1</sup>
$44/12$	ratio of molecular weights of CO <sub>2</sub> and carbon, dimensionless

$$G_{TOTAL,ij,t} = G_{w,ij,t} \cdot (1 + R_j) \quad (\text{B.4})$$

$$G_{w,ij,t} = I_{v,ij,t} \cdot D_j \cdot BEF_{1,j} \quad (\text{B.5})$$

where:

$G_{TOTAL,ij,t}$	average annual increment of total dry biomass of living trees for stratum i, species j, tonnes of dry matter, ha <sup>-1</sup> yr <sup>-1</sup> for year t
$G_{w,ij,t}$	average annual aboveground dry biomass increment of living trees for stratum i, species j, tonnes d.m. ha <sup>-1</sup> yr <sup>-1</sup> for year t
$R_j$	root-shoot ratio appropriate to increments for species j, dimensionless
$I_{v,ij,t}$	average annual increment in merchantable volume for stratum i, species j, m <sup>3</sup> ha <sup>-1</sup> yr <sup>-1</sup> for year t
$D_j$	basic wood density for species j, tonnes d.m. m <sup>-3</sup>
$BEF_{1,j}$	biomass expansion factor for conversion of annual net increment (including bark) in stem biomass to total aboveground tree biomass increment for species j dimensionless

**(b) Method 2 (stock change method)<sup>7</sup>**

$$\Delta C_{ij,t} = (C_{2,ij} - C_{1,ij}) / T_i \cdot 44/12 \quad (\text{B.6})$$

<sup>6</sup> GPG-LULUCF Equation 3.2.2, Equation 3.2.4 and Equation 3.2.5

<sup>7</sup> GPG-LULUCF Equation 3.2.3

$$C_{ij} = C_{AB,ij} + C_{BB,ij} \quad (\text{B.7})$$

$$C_{AB,ij} = A_{ij} \cdot V_{ij} \cdot D_j \cdot BEF_{2,j} \cdot CF_j \quad (\text{B.8})$$

$$C_{BB,ij} = C_{AB,ij} \cdot R_j \quad (\text{B.9})$$

where:

$\Delta C_{ij,t}$	average annual carbon stock change due to biomass growth of living trees for stratum $i$ , species $j$ , tonnes CO <sub>2</sub> yr <sup>-1</sup> for year $t$
$C_{2,ij}$	total carbon stock in living biomass of trees for stratum $i$ , species $j$ , calculated at time 2, tonnes C
$C_{1,ij}$	total carbon stock in living biomass of trees for stratum $i$ , species $j$ , calculated at time 1, tonnes C
$T_i$	number of years between times 2 and 1
$C_{AB,ij}$	carbon stock in aboveground tree biomass for stratum $i$ , species $j$ , tonnes C
$C_{BB,ij}$	carbon stock in belowground tree biomass for stratum $i$ , species $j$ , tonnes C
$A_{ij}$	area of stratum $i$ , species $j$ , hectare (ha)
$V_{ij}$	merchantable volume of stratum $i$ , species $j$ , m <sup>3</sup> ha <sup>-1</sup>
$D_j$	basic wood density for species $j$ , tonnes d.m. m <sup>-3</sup> merchantable volume
$BEF_{2,j}$	biomass expansion factor for conversion of stem biomass to aboveground tree biomass for species $j$ , dimensionless
$CF_j$	carbon fraction for species $j$ , tonnes C (tonne d.m.) <sup>-1</sup>
$R_j$	root-shoot ratio species $j$ , dimensionless

Time points 1 and 2, for which the stocks are estimated to determine  $\Delta C_{ij}$  must be broadly representative of the typical age of the trees under the baseline scenario during the crediting period. For example, if the trees are already mature at the start of the project, it is not appropriate to select time point 1 and 2 to correspond to the juvenile fast growth stage.

$C_{AB,ij}$  can alternatively be estimated through the use of an allometric equations and a growth model or yield table.

$$C_{AB,ij} = \sum_{l=1}^{N_{ij}} f_j(DBH_{ijl}, H_{ijl}) \cdot CF_j \cdot 0.001 \quad (\text{B.10})$$

where:

$C_{AB,ij}$	carbon stock in aboveground tree biomass for stratum $i$ , species $j$ , tonnes C
$N_{ij}$	number of trees of species $j$ in stratum $i$ , dimensionless
$f_j(DBH,H)$	allometric equation linking aboveground biomass of living trees (kg d.m. tree <sup>-1</sup> ) to mean diameter at breast height (DBH) and possibly tree height (H) for species $j$ .
$l$	sequence number of tree species $j$ in stratum $i$ , dimensionless
$CF_j$	carbon fraction for species $j$ , tonnes C (tonne d.m.) <sup>-1</sup>
0.001	conversion factor from kilograms to tonnes

For the choice of methods 1 or 2 above, there is no priority in terms of transparency and conservativeness. The choice should mainly depend on the kind of parameters available.  $V_{ij}$  and  $I_{v,ij}$  shall be estimated based on number of trees and national/local growth curve/table that usually can be obtained from national/local forestry inventory.  $D_j$ ,  $BEF_{1,j}$ ,  $BEF_{2,j}$ ,  $CF_j$  and  $R_j$  are regional and species specific and shall be chosen with the following priority:

- existing local and species specific;
- national and species specific (e.g. from national GHG inventory);

- c) species specific from neighbouring countries with similar conditions. In the case of a large country that encompasses very different biome types, c) might be preferable to b);
- d) globally species specific (e.g. GPG-LULUCF, IPCC 2006 Guideline for AFOLU).

If species specific information is unavailable, information for similar species (e.g., shape of trees, broadleaved vs. deciduous etc) can be used, with data source priority as listed for species specific information.

When choosing from global or national databases because local data are limited, it shall be confirmed with any available local data that this choice of values does not lead to underestimating the baseline net GHG removals by sinks, as far as can be judged. Local data used for confirmation may be drawn from the literatures and local forestry inventory or measured directly by project participants especially for BEF and root-shoot ratios that are age- and species- dependent.

Attention should be given to the fact that trees under the baseline scenario are trees outside forest and the biomass expansion factors (BEF<sub>2</sub>) for trees outside forest are generally higher than for forest trees. If BEF<sub>2</sub> from forests are used, the baseline net removals by sinks are subjected to be underestimated. Therefore, in case BEF<sub>2</sub> for trees outside forests are unavailable, to be conservative, the BEF<sub>2</sub> from forest trees shall be enlarged by 50%.

## 6. Additionality

This methodology uses the latest version of the “Tool for the demonstration and assessment of additionality for afforestation and reforestation CDM project activities” approved by the CDM Executive Board<sup>8</sup>, to demonstrate that a proposed A/R CDM project activity is additional and not the baseline scenario. Taking into account the conditions under which the proposed methodology is applicable, the tool is used, as follows:

**STEP 0:** *Preliminary screening based on the starting date of the project activity*

**STEP 1:** *Identification of alternatives to the project activity consistent with current laws and regulations*

**Sub-step 1a:** Define alternatives to the project activity

Explanation: As of conditions stated in section I.2 and demonstrated in section II.4 above, the lands to be afforested/reforested are severely degraded and the lands are still degrading. The lands are also economically unattractive. Apparent financial or other barriers prevent national and local governments from planting trees in the area of a proposed A/R CDM project activity. If these conditions are satisfied through analyses of socio-economic and environmental conditions (Section II.4 above and step 2 below), the third option, “continuation of the current situation” (no project activity or other alternatives undertaken) represents the only alternative to the proposed A/R CDM project activity. Other alternatives will be excluded in the proposed methodology.

**Sub-step 1b:** Enforcement of applicable laws and regulations

**STEP 2:** *Investment analysis*

**Sub-step 2a** Determine appropriate analysis method

Modification: Since the identified alternative (continuation of the current situation) does not need investments, the investment comparison analysis (Option II) will not be applicable in the proposed new methodology. Only Option I or Option III can be used for the proposed methodology.

---

<sup>8</sup> Throughout this document, “A/R additionality tool” refers to the document approved by the Executive Board of the CDM and available at the following URL: <http://cdm.unfccc.int/EB/Meetings/021/eb21repan16.pdf>.

**Sub-step 2b – Option I.** Apply simple cost analysis

**Sub-step 2b – Option II.** Investment comparison analysis

**Sub-step 2b – Option III.** Apply benchmark analysis

Notes: As an example, the PINFinancialAnalysis spreadsheet developed by the World Bank BioCarbon Fund<sup>9</sup> is a good template to do the benchmark analysis. The financial indicator of the template is FIRR and NPV with and without the CER benefit.

**Sub-step 2c** Calculation and comparison of financial indicators

Notes: More specifically for the A/R CDM project activity, the investment costs may include land purchase or rental, machinery, equipments, buildings, fences, site and soil preparation, seedling, planting, watering, weeding, pesticides, fertilization, supervision, training, technical consultation, etc. that occur in the establishment period. The operations and maintenance costs may include thinning, pruning, harvesting, replanting, fuel, transportation, repairs, fire and disease control, patrolling, administration, etc.) up to the end of crediting period. There are also transaction costs such as for project preparation, validation, registration, monitoring, etc. The revenues may include items like sale of timber, fuelwood, non-wood products and CER revenues.

**Sub-step 2d:** Sensitivity analysis

Notes: The most important drivers for uncertainties of the IRR/NPV may be the unit cost, and price of wood and non-wood products, because they are driven by future markets, and can be quite uncertain, especially in the longer term.

**STEP 3: Barrier analysis**

**Sub-step 3a:** Identify barriers that would prevent the implementation of the type of the proposed project activity

Modification: For an A/R CDM project activity, barriers may, in addition to those listed in the additionality tool, include:

- a) Investment barriers
  - ✓ The project owner cannot afford the high establishment investment in the early stage, because all economic returns, including from timber, non-wood products and CER, may occur ten or more years after the start of a proposed CDM project activity
  - ✓ No opportunity to get commercial loans from banks for the purpose of afforestation /reforestation because of the high market risk and economical unattractiveness in the context of arid and hyper arid deserts.
- b) Technological barriers, e.g.:
  - ✓ Lack of access to quality seed source and skills to produce high quality seedlings
  - ✓ Lack of skills for successful tree planting
  - ✓ Lack of skills for preventing planted trees from being subject to fire, pest and disease attack.
- c) Institutional barriers (e.g., lack of organizational instruments to integrate separate households and address technological barriers mentioned above)
  - ✓ High market risks

---

<sup>9</sup> [www.biocarfund.org](http://www.biocarfund.org)

**Sub-step 3b:** Show that the identified barriers would not prevent the implementation of at least one of the alternatives (except the proposed A/R CDM project activity already considered in step 3a)

Notes: The alternative land use (continued status as waste land) does, of course, not face the above-mentioned barriers.

**STEP 4: Impact of CDM registration**

**7. Ex ante actual net GHG removal by sinks**

**a. Verifiable changes in carbon stocks in the carbon pools**

The average annual carbon stock change in aboveground woody biomass, belowground woody biomass and soil organic matter between two monitoring events for stratum i, species j, species k can be estimated as described below.

$$\Delta C_{ijk,t} = (\Delta C_{AB,ijk,t} + \Delta C_{BB,ijk,t} + \Delta C_{SOC,ijk,t}) \cdot 44/12 \quad (\text{B.11})$$

where:

$\Delta C_{ijk,t}$	changes in carbon stock in carbon pools for stratum i, substratum j, species k, tonnes CO <sub>2</sub> yr <sup>-1</sup> for year t
$\Delta C_{AB,ijk,t}$	changes in carbon stock in aboveground woody biomass for stratum i, substratum j, species k, tonnes C yr <sup>-1</sup> for year t
$\Delta C_{BB,ijk,t}$	changes in carbon stock in belowground woody biomass for stratum i, substratum j, species k, tonnes C yr <sup>-1</sup> for year t
$\Delta C_{SOC,ijk,t}$	changes in carbon stock in soil organic matter for stratum i, substratum j, species k, tonnes C yr <sup>-1</sup> for year t
44/12	ratio of molecular weights of CO <sub>2</sub> and carbon, dimensionless

**a.1 Calculation of average annual carbon stock change in living woody biomass<sup>10</sup>**

$$\Delta C_{AB,ijk,t} = (C_{AB,t_2,ijk} - C_{AB,t_1,ijk}) / T_1 \quad (\text{B.12})$$

$$\Delta C_{BB,ijk,t} = (C_{BB,t_2,ijk} - C_{BB,t_1,ijk}) / T_1 \quad (\text{B.13})$$

$$C_{AB,ijk} = C_{AB\_tree,ijk} + C_{AB\_shrub,ijk} \quad (\text{B.14})$$

$$C_{BB,ijk} = C_{BB\_tree,ijk} + C_{BB\_shrub,ijk} \quad (\text{B.15})$$

Where

$\Delta C_{AB,ijk,t}$	changes in carbon stock in aboveground woody biomass for stratum i, substratum j, species k, tonnes C yr <sup>-1</sup> for year t
$\Delta C_{BB,ijk,t}$	changes in carbon stock in belowground woody biomass for stratum i, substratum j, species k, tonnes C yr <sup>-1</sup> for year t
$C_{AB,t_2,ijk}$	carbon stock in aboveground woody biomass for stratum i, substratum j, species k, calculated at time t <sub>2</sub> , tonnes C
$C_{AB,t_1,ijk}$	carbon stock in aboveground woody biomass for stratum i, substratum j, species k, calculated at time t <sub>1</sub> , tonnes C
$C_{BB,t_2,ijk}$	carbon stock in belowground woody biomass for stratum i, substratum j, species k, calculated at time t <sub>2</sub> , tonnes C

<sup>10</sup> Refers to equation 3.2.3 in GPG-LULUCF

$C_{BB, t1,ijk}$	carbon stock in belowground woody biomass for stratum i, substratum j, species k, calculated at time $t_1$ , tonnes C
$C_{AB\ tree,ijk}$	carbon stock in aboveground biomass of trees, tonnes C
$C_{AB\ shrub,ijk}$	carbon stock in aboveground biomass of planted shrubs, tonnes C
$C_{BB,tree,ijk}$	carbon stock in below-ground biomass of trees, tonnes C
$C_{BB\ shrub,ijk}$	carbon stock in below-ground biomass of planted shrubs, tonnes C
$T_1$	number of years between time $t_2$ and $t_1$ for biomass, $T_1 = t_2 - t_1$ , years

### a.1.1 Planted trees

The carbon stock in above- and below-ground biomass of pre-project existing trees shall not be included in the ex-ante estimation. Biomass Expansion Factors (BEF) method can be used to estimate the carbon stock in above- and below-ground biomass of living trees<sup>11</sup> that were planted within the A/R CDM project:

$$C_{AB\_tree,ijk} = A_{tree\_ijk} \cdot V_{tree\_ijk} \cdot D_k \cdot BEF_k \cdot CF_k \quad (\text{B.16})$$

$$C_{BB\_tree,ijk} = C_{AB\_tree,ijk} \cdot R_k \quad (\text{B.17})$$

where:

$C_{AB\ tree,ijk}$	carbon stock in aboveground biomass of trees, tonnes C
$C_{BB\ tree,ijk}$	carbon stock in belowground biomass of trees, tonnes C
$A_{tree,ijk}$	area covered by trees for stratum i, substratum j, species k, ha
$V_{tree\ ijk}$	mean merchantable/standing volume for stratum i, substratum j, and species k, $\text{m}^3 \text{ha}^{-1}$
$D_k$	volume-weighted average wood density for species k, tonnes d.m. $\text{m}^{-3}$
	merchantable/standing volume
$BEF_k$	biomass expansion factor for conversion of tree biomass of merchantable or standing volume to above-ground biomass, dimensionless
$CF_k$	carbon fraction, tonnes C (tonne d.m.) <sup>-1</sup> , IPCC default value = 0.5
$R_k$	root-shoot ratio, dimensionless

### a.1.2 Planted shrubs

$$C_{AB\_shrub,ijk} = A_{shrub,ijk} \cdot f(DB, H, C, N) \cdot CF_{s,k} \quad (\text{B.18})$$

$$C_{BB\_shrub,ijk} = C_{AB\_shrub,ijk} \cdot R_{s,k} \quad (\text{B.19})$$

where

$C_{AB\ shrub,ijk}$	carbon stock in aboveground biomass of shrubs, tonnes C
$C_{BB\ shrub,ijk}$	carbon stock in belowground biomass of shrubs, tonnes C
$A_{shrub,ijk}$	area of stratum i, substratum j, covered by shrub species k, hectare (ha)
$CF_{s,k}$	carbon fraction of shrub species k, dimensionless
$R_{s,k}$	root-shoot ratio of shrub species k, dimensionless
$f(DB, H, C, N)$	an allometric equation linking above-ground biomass (d.m. $\text{ha}^{-1}$ ) of shrubs to one or more of the variables diameter at base (DB), shrub height (H), crown area/diameter (C) and possibly number of stems (N)

The choice of methods and parameters shall be made in the same way as described in section II. 5. The changes of diameter at base (DB), shrub height (H), crown area/diameter (C) and number of stems (N) due to forage harvest and regrowth within each harvest cycle can be obtained from

<sup>11</sup> GPG-LULUCF Equation 3.2.3

literature, local yield table for planted shrubs, or surveys on similar shrub plantations that are under different stages of the harvest cycle in the vicinity. If there are no allometric equations available, or it is impossible to estimate the biomass of planted shrubs, the carbon stock change in living biomass of planted shrubs can be conservatively assumed to be zero.

### a.2. Calculation of average annual carbon stock change in soil organic matter<sup>12</sup>

The estimates of stock change in soil organic matter are based on the difference between an initial and final quasi-equilibrium (stable) soil C stock. The initial value is obtained from estimates for project lands before activities commence. The final value comes from a long-established forest stand of the same species, management practice and growing under conditions similar to those in the project area. A linear stock change is assumed to occur between the initial and the final soil C stock values, over a period of T years typically taken to reach the final soil C stock (IPCC default: 20 years). The stock change between initial and final states is divided by T as an estimate of the mean annual increment in mineral soil C under project conditions (see also Section 3.2.2.3.1.1 in GPG LULUCF).

$$\Delta SOC_{ijk,t} = (SOC_{For,ijk} - SOC_{Non-For,ij}) \cdot A_{ijk} / T_{For,ijk} \quad (\text{B.20})$$

Where:

$\Delta SOC_{ijk,t}$	average annual carbon stock change in soil organic matter for stratum i, substratum j, species k, tonnes C yr <sup>-1</sup>
$SOC_{For,ijk}$	stable soil organic carbon stock per hectare of plantation for stratum i, substratum j, species k, tonnes C ha <sup>-1</sup>
$SOC_{Non-For,ij}$	stable soil organic carbon stock per hectare of lands before planting for stratum i, substratum j, tonnes C ha <sup>-1</sup>
$A_{ijk}$	area of stratum i, substratum j, species k, hectare (ha)
$T_{For,ij}$	Duration of transition from $SOC_{Non-For,ij}$ to $SOC_{For,ijk}$ , year

$SOC_{For,ijk}$  should be preferably locally species- and management-specific and obtained from peer-reviewed scientific or other authoritative literatures or survey on plantations in the vicinity that better relate to species, rotation and management practices of planted forests under the project conditions, whenever possible.

If verifiable evidence (survey on similar plantations in vicinity or publications) indicates that the afforestation/reforestation would either increase or leave the soil organic carbon stock unchanged, the carbon stock changes in the soil organic matter can be conservatively assumed to be zero.

### b. GHG emissions by sources

$$GHG_{E,t} = E_{FossilFuel,t} + E_{Biomassloss,t} + N_2O_{N_{fixing},t} + N_2O_{Direct-N_{fertiliser},t} \quad (\text{B.21})$$

where

$GHG_{E,t}$	increase in GHG emission as a result of the implementation of the proposed A/R CDM project activity within the project boundary, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$E_{FossilFuel,t}$	increase in GHG emission as a result of burning of fossil fuels within the project boundary, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$E_{Biomassloss,t}$	CO <sub>2</sub> emissions as a result of a decrease in carbon stock in living biomass of existing non-tree vegetation, tonnes CO <sub>2</sub> . This is an initial loss, and therefore accounted once upfront as part of the first monitoring interval, not per year

<sup>12</sup> Refer to GPG-LULUCF Equation 3.2.31 and Equation 3.2.32

$N_2O_{N_{fixing},t}$  increase in N<sub>2</sub>O emission as a result of planting of N-fixing shrubs and cultivation of N-fixing annual crops within the project boundary, tonnes CO<sub>2</sub>-e yr<sup>-1</sup> for year t

$N_2O_{Direct-N_{fertiliser},t}$  increase in direct N<sub>2</sub>O emission as a result of nitrogen application within the project boundary, tonnes CO<sub>2</sub>-e.yr<sup>-1</sup> for year t

### b.1 Calculation of GHG emissions from burning fossil fuels

This most likely resulted from machinery use during site preparation and logging. IPCC 1996 Guideline could be used to estimate CO<sub>2</sub> emissions from combustion of fossil fuels:

$$E_{FossilFuel,t} = (CSP_{diesel,t} \cdot EF_{diesel} + CSP_{gasoline,t} \cdot EF_{gasoline}) \cdot 0.001 \quad (\text{B.22})$$

where:

$E_{FossilFuel,t}$  the increase in GHG emission as a result of burning of fossil fuels within the project boundary, tonnes CO<sub>2</sub>-e yr<sup>-1</sup> for year t

$CSP_{diesel,t}$  volume of diesel consumption in year t, litre (l) yr<sup>-1</sup>

$CSP_{gasoline,t}$  volume of gasoline consumption in year t, litre (l) yr<sup>-1</sup>

$EF_{diesel}$  emission factor for diesel, kg CO<sub>2</sub> l<sup>-1</sup>

$EF_{gasoline}$  emission factor for gasoline, kg CO<sub>2</sub> l<sup>-1</sup>

0.001 conversion from kg to tonnes of CO<sub>2</sub>

Project participants should use national CO<sub>2</sub> emission factors. If these are unavailable they may use default emission factors as provided in the 1996 Revised IPCC Guidelines, GPG 2000 and IPCC 2006 Guidelines.

### b.2 Calculation of the decrease in carbon stock in living biomass of existing non-tree vegetation

It is assumed that all existing non-tree vegetation will disappear due to site preparation or competition from planted trees. The carbon stock in existing non-tree vegetation can be estimated using local or national data available or by direct measurement during baseline survey using method described in Section III.5.b.2 below. This is a conservative assumption because there will be some non-tree vegetation in the project scenario. Some vegetation may re-grow even if all non-tree vegetation is removed during the site preparation (overall plough). The non-tree vegetation carbon loss will be accounted once during the crediting period, as part of the first monitoring interval.

$$E_{biomassloss,t} = \sum_i A_i \cdot B_{non-tree,i} \cdot CF_{non-tree} \cdot 44/12 \quad (\text{B.23})$$

where:

$E_{Biomassloss,t}$  CO<sub>2</sub> emissions as a result of a decrease in carbon stock in living biomass of existing non-tree vegetation, tonnes CO<sub>2</sub>. This is an initial loss, and therefore accounted once upfront as part of the first monitoring interval, not per year

$A_i$  area of stratum i, ha

$B_{non-tree,i}$  average non-tree biomass stock on land to be planted before the start of a proposed A/R CDM project activity for stratum i, tonnes d.m. ha<sup>-1</sup>

$CF_{non-tree}$  carbon fraction of dry biomass in non-tree vegetation, tonnes C (tonne d.m.)<sup>-1</sup>

44/12 ratio of molecular weights of CO<sub>2</sub> and carbon, dimensionless



**b.3 Calculation of nitrous oxide emissions from nitrogen fertilization practices<sup>1314</sup>**

$$N_2O_{direct-N_{fertilizer,t}} = [(F_{SN,t} + F_{ON,t}) \cdot EF_1] \cdot 44/28 \cdot GWP_{N_2O} \quad (B.24)$$

$$F_{SN,t} = N_{SN-Fert,t} \cdot (1 - Frac_{GASF}) \quad (B.25)$$

$$F_{ON,t} = N_{ON-Fert,t} \cdot (1 - Frac_{GASM}) \quad (B.26)$$

where:

$N_2O_{direct-N_{fertilizer,t}}$	direct N <sub>2</sub> O emission as a result of nitrogen application within the project boundary, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$F_{SN,t}$	mass of synthetic fertilizer nitrogen applied adjusted for volatilization as NH <sub>3</sub> and NO <sub>x</sub> , tonnes N yr <sup>-1</sup> in year t
$F_{ON,t}$	annual mass of organic fertilizer nitrogen applied adjusted for volatilization as NH <sub>3</sub> and NO <sub>x</sub> , tonnes N yr <sup>-1</sup> in year t
$N_{SN-Fert,t}$	mass of synthetic fertilizer nitrogen applied, tonnes N yr <sup>-1</sup> in year t
$N_{ON-Fert,t}$	mass of organic fertilizer nitrogen applied, tonnes N yr <sup>-1</sup> in year t
$EF_1$	emission factor for emissions from N inputs, tonnes N <sub>2</sub> O-N (tonnes N input) <sup>-1</sup>
$Frac_{GASF}$	the fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for synthetic fertilizers, tonnes NH <sub>3</sub> -N and NO <sub>x</sub> -N (tonnes N input) <sup>-1</sup>
$Frac_{GASM}$	the fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for organic fertilizers, tonnes NH <sub>3</sub> -N and NO <sub>x</sub> -N (tonnes N input) <sup>-1</sup>
44/28	ratio of molecular weights of N <sub>2</sub> O and nitrogen, dimensionless
$GWP_{N_2O}$	global warming potential for N <sub>2</sub> O (with a value of 310 for the first commitment period)

As noted in GPG 2000, the default emission factor (EF<sub>1</sub>) is 1.25 % of applied N, and this value should be used when country-specific factors are unavailable. The default values for the fractions of synthetic and organic fertiliser nitrogen that are emitted as NO<sub>x</sub> and NH<sub>3</sub> are 0.1 and 0.2 respectively in 1996 IPCC Guideline. Project participants may use scientifically-established specific emission factors that are more appropriate for their project. Specific good practice guidance on how to derive specific emission factors is given in Box 4.1 of GPG 2000. Updated values shall be used upon the publication of IPCC 2006 Guidelines.

**b.4 GHG emissions from intercropping of N-fixing shrub species and of N-fixing annual crop<sup>15</sup>**

$$N_2O_{N_{fixing,t}} = (F_{BN,t} + F_{SN,t}) \cdot EF_1 \cdot 44 / 28 \cdot GWP_{N_2O} \quad (B.27)$$

$$F_{BN,t} = \sum_i \sum_j \sum_k (Crop_{BF_k,t} \cdot Crop_{RA_k} \cdot Crop_{NCRBF_k} \cdot A_{ijk}) \quad (B.28)$$

$$F_{SBN,t} = \sum_i \sum_j \sum_k (\Delta B_{AB\_shrub_{ijk,t}} \cdot LF_k \cdot Shrub_{NCRBF_k} \cdot A_{ijk}) \quad (B.29)$$

<sup>13</sup> Refers to Equation 3.2.18 in IPCC GPG-LULUCF, Equation 4.22, Equation 4.23 and Equation 4.3.1 in GPG-2000

<sup>14</sup> Based on the guidance on accounting for emissions of N<sub>2</sub>O from fertilizer application agreed on EB 26<sup>th</sup> meeting, only direct (e.g. volatilization) emissions of N<sub>2</sub>O from application of fertilizers within the project boundary shall be accounted for in A/R project activities.

<sup>15</sup> Refers to Equation 4.20 and Equation 4.25-4.27 in IPCC GPG-2000 for Agriculture.

where

$N_2O_{N_{fixing},t}$	increase in annual N <sub>2</sub> O emission as a result of planting of N-fix shrubs and intercropping of N-fixing annual crop within the project boundary, tonnes CO <sub>2</sub> -e. yr <sup>-1</sup> in year t
$F_{BN,t}$	amount of nitrogen fixed by N-fixing intercrops cultivated annually, t N yr <sup>-1</sup>
$F_{SBN,t}$	amount of nitrogen fixed by N-fixing shrubs planted, t N. yr <sup>-1</sup> in year t
$EF_1$	emission factor for emissions from N inputs, tonnes N <sub>2</sub> O-N (tonnes N input) <sup>-1</sup>
$Crop_{BF_k,t}$	the seed yield of N-fixing intercrops per hectare for crop type k, t d.m. ha <sup>-1</sup> yr <sup>-1</sup> in year t
$Crop_{RA_k,t}$	the ratio of dry matter in the aboveground crop biomass (including residue) to the seed yield for crop type k, dimensionless
$Crop_{NCRBF_k}$	the fraction of crop biomass that is nitrogen for crop type k, dimensionless
$A_{ijk}$	area of N-fixing intercrops or shrubs for stratum i, substratum j, crop type or species k, ha
$\Delta B_{AB\_shrub_{ijk,t}}$	annual stock change of aboveground biomass for stratum i, substratum j, shrub species k, t d.m. ha <sup>-1</sup> yr <sup>-1</sup> in year t
$LF_k$	the ratio of leaf biomass in aboveground biomass of N-fixing shrubs, dimensionless
$Shrub_{NCRBF_k}$	the fraction of N-fixing shrub biomass that is nitrogen for species k, dimensionless
$GWP_{N_2O}$	global warming potential for N <sub>2</sub> O (with a value of 310 for the first commitment period)

Country or local specific value for  $Crop_{RA_k}$ ,  $Crop_{NCRBF_k}$  and  $Shrub_{NCRBF_k}$  shall be used. If country-specific data are not available, the default values may be chosen from Table 4.16 of GPG-2000, Table 4-19 of the Reference Manual of the IPCC 1996 Guidelines (0.03 kg N (kg dry matter)<sup>-1</sup>) and related tables in IPCC 2006 Guidelines.

Country-specific emission factor (EF<sub>1</sub>) should be used where possible, in order to reflect the specific conditions of a country and the agricultural practices involved. If country-specific emission factor is not available, EF<sub>1</sub> from other countries with comparable management and climatic conditions are good alternatives. Otherwise, the default emission factor (EF<sub>1</sub>) is 1.25 % of input N as noted in GPG-2000 or updated value in IPCC 2006 Guidelines. There is no default value for LF<sub>k</sub>, so it may be found in literatures, otherwise it should be obtained through direct measurement.

### c. Actual net GHG removals by sinks

The actual net greenhouse gas removals are calculated as follows:

$$C_{ACTUAL,t} = \sum_i \sum_j \sum_k \Delta C_{ijk,t} - GHG_{E,t} \quad (\text{B.30})$$

where:

$C_{ACTUAL,t}$	Actual net greenhouse gas removals by sinks, tonnes CO <sub>2</sub> -e yr <sup>-1</sup>
$\Delta C_{ijk,t}$	average annual carbon stock change in living biomass of trees for stratum i, species j, tonnes CO <sub>2</sub> yr <sup>-1</sup> for year t
$GHG_{E,t}$	GHG emissions by sources within the project boundary as a result of the implementation of an A/R CDM project activity, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t

## 8. Leakage

Leakage represents the increase in GHG emissions by sources, which occurs outside the boundary of an A/R CDM project activity, and which is measurable and attributable to the A/R CDM project activity. As per the applicability conditions, the proposed A/R CDM project activity will provide at least the same amount of goods and services for local communities as in the absence of the project

activity, and there is no grazing in either the project scenario or baseline scenario. As a result, leakage due to shift of pre-project activities is nil. However, to increase the income of local communities or to improve the financial revenue stream of the project, some projects may intentionally plant forage species among tree rows. The production of forage will support the raising of livestock outside the project boundary, and as a result will increase leakage emissions due to enteric fermentation and manure management outside the project boundary.

In addition, in the context of A/R activities, there will be leakage GHG emissions from fossil fuel combustion from vehicle use due to the transportation of seedlings, workers, staff, and harvest products, to and/or from the project sites. Emission sources included in or excluded from leakage in the proposed methodology are listed in Table 3 below.

**Table 3: Emissions sources included in or excluded from leakage.**

Source	Gas	Included/ excluded	Justification / Explanation of choice
Combustion of fossil fuels	CO <sub>2</sub>	Included	Potential significant emission source
	CH <sub>4</sub>	Excluded	Potential emission is negligibly small
	N <sub>2</sub> O	Excluded	Potential emission is negligibly small
Activity displacement (grazing, fuelwood collecting)	CO <sub>2</sub>	Excluded	As per the applicability condition of the proposed methodology, there is no grazing either in the project case or the baseline scenario, and the project can provide the same amount of goods and services as in the absence of the A/R CDM project activity
	CH <sub>4</sub>	Excluded	
	N <sub>2</sub> O	Excluded	
Use of fertilizers	CO <sub>2</sub>	Excluded	Not applicable
	CH <sub>4</sub>	Excluded	Not applicable
	N <sub>2</sub> O	Excluded	Potential emission from N-fertilizer application in nursery is negligibly small as per EB26 decision
Livestock fed with forage produced by the project	CO <sub>2</sub>	Excluded	Not applicable
	CH <sub>4</sub>	Included	Potential significant emission source
	N <sub>2</sub> O	Included	Potential significant emission source

There are two sources of the leakage covered by this methodology:

- GHG emissions caused by vehicle fossil fuel combustion due to transportation of seedlings, labour, staff and harvest products to or from project sites (while avoiding double-counting with emissions accounted for in  $E_{\text{FossilFuel}}$  above).
- GHG emissions from livestock fed with forage produced by the project activities (forage-fed livestock).

$$LK_t = LK_{\text{Vehicle}, \text{CO}_2, t} + LK_{\text{FFL}, t} \quad (\text{B.31})$$

where:

- $LK_t$  Leakage due to the increase in GHG emissions by sources outside the project boundary and attributable to the A/R CDM project activity, tonnes CO<sub>2</sub>-e yr<sup>-1</sup> for year t
- $LK_{\text{Vehicle}, \text{CO}_2, t}$  GHG emissions due to fossil fuel combustion from vehicles, tonnes CO<sub>2</sub>-e yr<sup>-1</sup> for year t
- $LK_{\text{FFL}, t}$  GHG emissions from the forage-fed livestock, tonnes CO<sub>2</sub>-e yr<sup>-1</sup> for year t

**a. Estimation of  $LK_{\text{vehicle}}$  (leakage due to fossil fuel consumption)**

The CO<sub>2</sub> emissions due to fossil fuel consumption of vehicles can be estimated using the bottom-up approach described in the GPG 2000<sup>16</sup>.

$$LK_{\text{Vehicle}, \text{CO}_2, t} = \sum_i \sum_j (EF_{ij} \cdot \text{FuelConsumption}_{ij, t}) / 1000 \quad (\text{B.32})$$

$$\text{FuelConsumption}_{ij, t} = n_{ij, t} \cdot k_{ij, t} \cdot e_{ij} \quad (\text{B.33})$$

where:

$LK_{\text{Vehicle}, \text{CO}_2, t}$	Total CO <sub>2</sub> emissions due to fossil fuel combustion from vehicles, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>
<i>i</i>	Vehicle type
<i>j</i>	Fuel type
$EF_{ij}$	Emission factor for vehicle type <i>i</i> with fuel type <i>j</i> , kg CO <sub>2</sub> /litre
$\text{FuelConsumption}_{ij}$	Consumption of fuel type <i>j</i> of vehicle type <i>i</i> , litres for year <i>t</i>
$n_{ij, t}$	Number of vehicles for year <i>t</i>
$k_{ij, t}$	Kilometres travelled by each of vehicle type <i>i</i> with fuel type <i>j</i> , km for year <i>t</i>
$e_{ij}$	Average fuel consumption of vehicle type <i>i</i> with fuel type <i>j</i> , litres km <sup>-1</sup>

Country-specific emission factors shall be used if available. Default emission factors provided in the IPCC Guidelines and updated in the GPG 2000 and IPCC 2006 Guidelines may be used if there are no country-specific data.

**b. Estimation of  $LK_{\text{FFL}}$  (leakage from forage-fed livestock)**

The following types of leakage of GHG emissions from forage-fed livestock are accounted for:

- CH<sub>4</sub> emissions from enteric fermentation by the forage-fed livestock;
- CH<sub>4</sub> emissions from manure management for the forage-fed livestock;
- N<sub>2</sub>O emissions from manure management for the forage-fed livestock

$$LK_{\text{FFL}, t} = LK_{\text{CH}_4, \text{FFL}, \text{Ferm}, t} + LK_{\text{CH}_4, \text{FFL}, \text{manure}, t} + LK_{\text{N}_2\text{O}, \text{FFL}, \text{manure}, t} \quad (\text{B.34})$$

where:

$LK_{\text{FFL}, t}$	GHG emissions from the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>
$LK_{\text{CH}_4, \text{FFL}, \text{Ferm}, t}$	CH <sub>4</sub> emissions from enteric fermentation by the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>
$LK_{\text{CH}_4, \text{FFL}, \text{manure}, t}$	CH <sub>4</sub> emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>
$LK_{\text{N}_2\text{O}, \text{FFL}, \text{manure}, t}$	N <sub>2</sub> O emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>

As specified as an applicability condition, all forage produced by the project shall have a similar nutritional value and digestibility, and will support only a single livestock group with a single manure management system. If these conditions are not met, this methodology can not be used.

For *ex ante* estimates of leakage emissions, a suitable livestock group and manure management system may be specified according to knowledge of intended project activities, forage types, and local farming practices<sup>17</sup>. Alternatively, the forage-fed livestock group may be selected *ex ante* as the pre-

<sup>16</sup> Refer to Equation 2.5 and Equation 2.6 in IPCC GPG 2000 for energy sector

<sup>17</sup> In this case, only this livestock group and manure management system can be used by farmers that utilise forage provided by the project.

project livestock group that is fed the largest amount of forage most similar to that to be produced by the project, as determined from data obtained by a survey on livestock forage feeding from households likely to be involved in the project—at least 30 households or 10% of households, whichever is greater, should be sampled. The manure management system to be used in *ex ante* emissions estimation shall be selected as the most common manure management system for the identified forage-fed livestock group. Characteristics of the forage-fed livestock group that will help select appropriate enteric CH<sub>4</sub> emission factors should also be identified and determined—by household survey if necessary—including, for example, mean weight, growth rate, and milk production.

### b.1 CH<sub>4</sub> emissions from enteric fermentation of forage-fed livestock ( $LK_{CH_4_{FFL,Ferm,t}}$ )

CH<sub>4</sub> emissions from enteric fermentation by the forage-fed livestock can be estimated based on forage production, daily biomass intake of the fed animals, and emission factors as per IPCC GPG 2000 and IPCC 2006 Guidelines for AFOLU, using the equation below<sup>18</sup>

$$LK_{CH_4_{FFL,Ferm,t}} = EF_1 \cdot Population_t \cdot 0.001 \cdot GWP_{CH_4} \quad (B.35)$$

$$Population_t = Produc_{Forage,t} / (DBI \cdot 365) \quad (B.36)$$

where:

$LK_{CH_4_{FFL,Ferm,t}}$	CH <sub>4</sub> emissions from enteric fermentation by the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$EF_1$	Enteric CH <sub>4</sub> emission factor for the forage-fed livestock, kg CH <sub>4</sub> head <sup>-1</sup> yr <sup>-1</sup>
$Population_t$	Equivalent number of forage-fed livestock, head for year t
$Produc_{Forage,t}$	Production of forage by the project in year t, kg d.m. yr <sup>-1</sup>
$DBI$	Daily biomass intake for the forage-fed livestock, kg d.m. head <sup>-1</sup> day <sup>-1</sup>
$GWP_{CH_4}$	Global warming potential for CH <sub>4</sub> (with a value of 23 for the first commitment period)
0.001	Conversion of kilograms into tonnes, dimensionless
365	Number of days per year

The production of forage can be estimated by collecting production rates from literature that represents the shrub species, climate, soil conditions and other features of the area in which the forage will be grown. The production rate can also be estimated by sampling surveys of forage crops in the vicinity that produce the same type of forage in areas with similar soil conditions.

Country-specific emission factors for enteric CH<sub>4</sub> emissions, which have been fully documented in peer reviewed publications or are from national GHG inventory, are preferable. Otherwise, methane emission factors can be taken from Table 10.10 and Table 10.11 in the IPCC 2006 Guidelines for AFOLU. When selecting emission factors it is important to select those from a region similar to the project area. Also, scrutinise the tabulations in Annex 10A.1 of the IPCC 2006 Guideline for AFOLU to ensure that the underlying animal characteristics such as weight, growth rate and milk production used to develop the emission factors are similar to those attained for local conditions. In particular, data collected on average annual milk production by dairy cows should be used to help select a dairy cow emission factor. Data that have been fully documented in peer reviewed publications, or are from national GHG inventory, may also be used. If necessary, interpolate between dairy cow emission factors shown, for example, in IPCC Annex 10A.1, using the data collected on average annual milk production per head during the household survey.

<sup>18</sup> Refers to equation 10.19 and equation 10.20 in AFLOU volume of the IPCC 2006 Guidelines or equation 4.12 and equation 4.13 in GPG 2000 for agriculture

For data on daily biomass intake, preferably use local data, or applicable data that have been fully documented in peer reviewed publications or are from national GHG inventory. When selecting the value of daily biomass intake, ensure that the chosen data are applicable to both the forage types to be produced by the project and the livestock group to be supported by the forage. For examples of default data on daily biomass intake by livestock group, see Table 4 below.

**Table4: Approximate values of daily biomass intake for different livestock groups<sup>19</sup>**

Livestock groups	Country Group	Daily Feed Intake (MJ head <sup>-1</sup> day <sup>-1</sup> )	Daily Biomass Intake (kg d.m. head <sup>-1</sup> day <sup>-1</sup> )
Sheep	Developed Countries	20	2.0
	Developing Countries	13	1.3
Goats	Developed Countries	14	1.4
	Developing Countries	14	1.4
Mules / Asses	Developed Countries	60	6.0
	Developing Countries	60	6.0

## b.2 CH<sub>4</sub> emissions from manure management for forage-fed livestock ( $LK_{CH_{4FFLmanure,t}}$ )

The storage and treatment of manure under anaerobic conditions will produce CH<sub>4</sub>. These conditions occur most readily when large numbers of animals are managed in a confined area (e.g., dairy farms, beef feedlots, and swine and poultry farms), and where manure is disposed of in liquid-based systems. The main factors affecting CH<sub>4</sub> emissions are the amount of manure produced and the portion of the manure that decomposes anaerobically. The former depends on the rate of waste production per animal and the number of animals, and the latter on how the manure is managed. When manure is stored or treated as a liquid (e.g. in lagoons, ponds, tanks, or pits), it decomposes anaerobically and can produce a significant quantity of CH<sub>4</sub>. The temperature and the retention time of storage greatly affect the amount of methane produced. When manure is handled as a solid (e.g. in stacks or piles), or when it is deposited on pastures and rangelands, it tends to decompose under more aerobic conditions and less CH<sub>4</sub> is produced.

CH<sub>4</sub> emissions from manure management for the forage-fed livestock can be estimated using IPCC methods<sup>20</sup>:

$$LK_{CH_{4FFL,manure,t}} = EF_2 \cdot Population_t \cdot 0.001 \cdot GWP_{CH_4} \quad (B.37)$$

where

$LK_{CH_{4FFL,manure,t}}$	CH <sub>4</sub> emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$Population_t$	Equivalent number of forage-fed livestock supported by the project, head for year t
$EF_2$	Manure management CH <sub>4</sub> emission factor for the forage-fed livestock, kg CH <sub>4</sub> head <sup>-1</sup> yr <sup>-1</sup>
$GWP_{CH_4}$	Global warming potential for CH <sub>4</sub> (with a value of 23 for the first commitment period)
0.001	Conversion of kilograms into tonnes, dimensionless

The best estimates of emissions will usually be obtained using country-specific emission factors that have been fully documented in peer-reviewed publications or are from national GHG inventory. It is

<sup>19</sup> Data from AR-AM0003/version 2.

<sup>20</sup> Refers to equation 10.22 in AFLOU volume of the IPCC 2006 Guidelines or equation 4.15 in GPG 2000 for agriculture

recommended that country-specific emission factors be used that reflect the actual duration of storage and type of treatment of animal manure in the management system used. If appropriate country-specific emission factors are unavailable, default emission factors presented in Table 10.14–10.16 of the IPCC 2006 Guidelines for AFOLU may be used. These emission factors represent those for a range of livestock types and associated manure management systems, by regional management practice and temperature. When selecting a default factor, be sure to consult the supporting tables in Annex 10A.2 of IPCC 2006 Guidelines for AFOLU, for the distribution of manure management systems and animal waste characteristics used to estimate emissions. Select an emission factor for a region that most closely matches the circumstances of the livestock that are fed forage from the project.

### b.3 N<sub>2</sub>O emissions from manure management for forage-fed livestock ( $LK_{N_2O_{FFL,manure,t}}$ )

Nitrous oxide emissions from manure management vary significantly between the type of management system used, and can also result in indirect emissions due to other forms of nitrogen loss from the system. As specified by an applicability condition, forage produced by the project must support only one livestock group and the group must have a single manure management system. The N<sub>2</sub>O emissions from manure management for the forage-fed livestock supported by the project can be estimated using methods provided in the IPCC 2006 Guidelines for AFOLU, or in IPCC GPG 2000<sup>21</sup>:

$$LK_{N_2O_{FFL,manure,t}} = LK_{Direct\_N_2O,manure,t} + LK_{Indirect\_N_2O,manure,t} \quad (\text{B.38})$$

$$LK_{Direct\_N_2O_{FFL,manure,t}} = Population_t \cdot Nex \cdot EF_3 \cdot 0.001 \cdot 44 / 28 \cdot GWP_{N_2O} \quad (\text{B.39})$$

$$LK_{Indirect\_N_2O_{FFL,manure,t}} = Population_t \cdot Nex \cdot Frac_{Gas} \cdot EF_4 \cdot 0.001 \cdot 44 / 28 \cdot GWP_{N_2O} \quad (\text{B.40})$$

where

$LK_{N_2O_{FFL,manure,t}}$	N <sub>2</sub> O emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$LK_{Direct\_N_2O_{FFL,manure,t}}$	Direct N <sub>2</sub> O emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$LK_{Indirect\_N_2O_{FFL,manure,t}}$	Indirect N <sub>2</sub> O emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
Population <sub>t</sub>	Equivalent number of forage-fed livestock, head, for year t
Nex	Annual average N excretion per head for the forage-fed livestock, kg N head <sup>-1</sup> yr <sup>-1</sup>
EF <sub>3</sub>	Emission factor for direct N <sub>2</sub> O emission from manure management for the forage-fed livestock, kg N <sub>2</sub> O-N (kg N) <sup>-1</sup>
EF <sub>4</sub>	Emission factor for N <sub>2</sub> O emissions from atmospheric deposition of forage-sourced nitrogen on soils and water surfaces, kg N <sub>2</sub> O-N (kg NH <sub>3</sub> -N and NO <sub>x</sub> -N emitted) <sup>-1</sup> . Use of the IPCC default factor of 0.01 is recommended.
Frac <sub>Gas</sub>	Fraction of managed livestock manure nitrogen that volatilises as NH <sub>3</sub> and NO <sub>x</sub> in the manure management phase for the forage-fed livestock, kg NH <sub>3</sub> -N and NO <sub>x</sub> -N (kg N) <sup>-1</sup>
GWP <sub>N<sub>2</sub>O</sub>	Global Warming Potential for N <sub>2</sub> O (= 310 in the 1 <sup>st</sup> C.P.)
44/28	Conversion of (N <sub>2</sub> O-N) emissions to N <sub>2</sub> O emissions, dimensionless
0.001	Conversion of kilograms to tonnes, dimensionless

Annual nitrogen excretion rates must be determined for the forage-fed livestock group supported by

<sup>21</sup> Refers to equations 10.25, 10.26 and 10.27 in AFLOU volume of the IPCC 2006 Guidelines and/or equation 4.18 in GPG 2000 for agriculture

the project. The best estimates of excretion rates will usually be obtained using country-specific rates that have been fully documented in peer-reviewed publications or are from national GHG inventory. If country-specific data cannot be collected or derived, or appropriate data are not available from another country, default nitrogen excretion rates presented in Table 10.19 of IPCC 2006 Guidelines for AFOLU can be used.

The best estimates of emissions will usually be obtained using country-specific emission factors that have been fully documented in peer-reviewed publications or are from national GHG inventory. It is recommended to use country-specific emission factors that reflect the actual duration of storage and type of treatment of animal manure in the management system that is used. If appropriate country-specific emission factors are unavailable, the default emission factors presented in Table 10.21 and Table 11.3 of the IPCC 2006 Guidelines for AFOLU can be used. Default values for volatilization of  $\text{NH}_3$  and  $\text{NO}_x$  ( $\text{Frac}_{\text{Gas}}$ ) in the manure management system are presented in the Table 10.22 of the IPCC 2006 Guidelines.

The default value for  $\text{EF}_4$  in equation 10.27 of the IPCC 2006 Guidelines for AFOLU is 0.01 tonnes  $\text{N}_2\text{O-N}$  (tonnes  $\text{NH}_3\text{-N}$  and  $\text{NO}_x\text{-N}$  emitted)<sup>-1</sup>. Country-specific values for  $\text{EF}_4$  should be used with great care because of the special complexity of transboundary atmospheric transport: instead, use of IPCC default values is recommended. This is because although specific countries may have specific measurements of N deposition and associated  $\text{N}_2\text{O}$  flux, in many cases the deposited N may not have originated in their country. Similarly, some of the N that volatilises in their country may be transported to and deposited in another country, where different conditions that affect the fraction emitted as  $\text{N}_2\text{O}$  may prevail.

## 9. Ex ante net anthropogenic GHG removal by sinks

The following general formula can be used to calculate the net anthropogenic GHG removals by sinks of an A/R CDM project activity ( $C_{\text{AR-CDM},t}$ ), in tonnes  $\text{CO}_2\text{-e yr}^{-1}$ :

$$C_{\text{AR-CDM},t} = C_{\text{ACTUAL},t} - C_{\text{BSL},t} - LK_t \quad (\text{B.41})$$

where:

$C_{\text{AR-CDM},t}$	net anthropogenic GHG removal by sink for year t, tonnes $\text{CO}_2\text{-e yr}^{-1}$
$C_{\text{ACTUAL},t}$	actual net anthropogenic GHG removal by sink for year t, tonnes $\text{CO}_2\text{-e yr}^{-1}$
$C_{\text{BSL},t}$	baseline net greenhouse gas removals by sinks for year t, tonnes $\text{CO}_2\text{-e yr}^{-1}$
$LK_t$	leakage, $\text{CO}_2\text{-e yr}^{-1}$ for year t

Please see section III.9 for the formulae to calculate net anthropogenic GHG removals by sinks for project activities using tCERs and for those using ICERs, which is based on EB 22 annex 15.

## 10. Uncertainties

### a. Uncertainties to be considered

This methodology uses methods from IPCC GPG-LULUCF, GPG 2000, IPCC 2006 Guidelines as well as related rules for A/R CDM project activities to estimate the baseline net GHG removals by sinks, the leakage, the actual net GHG removals by sinks and the net anthropogenic GHG removals by sinks. Potential uncertainties arise from emission factors and sampling surveys. These uncertainties and their countermeasures are elaborated below.

- Uncertainties arising from, for example, biomass expansion factors (BEFs) or basic wood density would result in uncertainties in the estimation of both the baseline net GHG removals by sinks and the actual net GHG removals by sinks, especially when global default values are used. This methodology recommends project participants to identify key parameters that would significantly influence the estimation results, and to try to develop local values for key



factors using various data sources including direct measurement, or to choose conservative values.

- b) Uncertainties arising from sample survey (statistical uncertainties): The sampling error for each stratum may result from large spatial variability. Therefore an appropriate sampling protocol is necessary, including sufficient number of samples, variation and uncertainty analysis, sound quality control and quality assurance.

### b. Uncertainty assessment

The percentage uncertainty on the estimate of certain parameters and data (yield table values, biomass expansion factors, wood density, carbon fraction and other biophysical parameters) can be assessed from the sample standard deviation of measured sample values, using half the 95% confidence interval width divided by the estimated value, i.e.<sup>22</sup>,

$$U_s (\%) = \frac{1/2(95\% \text{ConfidenceIntervalWidth})}{\mu} \cdot 100 \quad (\text{B.42})$$

$$= \frac{1/2(4\sigma)}{\mu} \cdot 100$$

where

- $U_s$  = percentage uncertainty on the estimate of the mean parameter value, %
- $\mu$  = sample mean value of the parameter
- $\sigma$  = sample standard deviation of the parameter

If the default parameters are used, uncertainty will be higher than if locally measured parameters are used, and can be only roughly estimated with expert judgment<sup>23</sup>.

The percentage uncertainties on quantities that are the product of several terms are then estimated using the following equation<sup>24</sup>:

$$U_s = \sqrt{U_1^2 + U_2^2 + \dots + U_n^2} \quad (\text{B.43})$$

where

- $U_s$  = percentage uncertainty of product (emission by sources or removal by sinks);
- $U_i$  = percentage uncertainties associated with each term of the product (parameters and activity data),  $i=1,2,\dots,n$

The percentage uncertainty on quantities that are the sum or difference of several terms can be estimated using following simple error propagation equation<sup>25</sup>:

$$U_c = \frac{\sqrt{(U_{s1} \cdot C_{s1})^2 + (U_{s2} \cdot C_{s2})^2 + \dots + (U_{sn} \cdot C_{sn})^2}}{|C_{s1} + C_{s2} + \dots + C_{sn}|} \quad (\text{B.44})$$

where

- $U_c$  = combined percentage uncertainty, %
- $U_{si}$  = percentage uncertainty on each term of the sum or difference, %
- $C_{si}$  = mean value of each term of the sum or difference

<sup>22</sup> Box 5.2.1 in GPG LULUCF

<sup>23</sup> GPG LULUCF Chapter 5.2 and Chapter 3.2

<sup>24</sup> Equation 5.2.1 in GPG LULUCF

<sup>25</sup> Refers to equation 5.2.2 in GPG LULUCF

**c. Countermeasure for Reducing Statistical Uncertainties**

This methodology can basically reduce uncertainties through:

- a) Proper stratification of the project area into relatively homogeneous strata;
- b) Conservativeness in the selection of carbon pools and in setting values for BEFs and root-shoot ratios.

**d. Conservativeness will be achieved through**

- The baseline methodology set the zero carbon stock change for lands without growing trees and projects the continuous biomass growth for lands with growing trees. This simple baseline methodology ensures the baseline scenario is conservative.
- The BEF values for pre-project trees are conservatively set as 150% of those for forest trees.
- The omitting of the pools of dead wood and litter is a conservative way.
- The methodology that accounts all pre-existing non-tree vegetation as an emission, which is conservative.

**11. Data needed for ex ante estimations**

**Table 5: Data/parameter, their vintage, geographical scale and possible data sources**

<b>Data / parameter</b>	<b>Description</b>	<b>Vintage</b>	<b>Data sources and geographical scale</b>
Historical land use/cover data	Determining baseline approach Demonstrating eligibility of land	Earliest possible up to now	Publications, national or regional forestry inventory, local government, interview, PRA survey
Land use/cover map	Demonstrating eligibility of land, stratifying land area	Around 1990 and most recent date	National, regional or local forestry inventory
Satellite image	Demonstrating eligibility of land, stratifying land area	1989/1990 and most recent date	e.g. local Landsat
Landform map	Stratifying land area	most recent date	1:100,000 or higher from local government and institutional agencies
Soil map	Stratifying land area	most recent date	1:100,000 or higher from local government and institutional agencies
National and sectoral policies	Additionality consideration	since 1998	National and sectoral
UNFCCC decisions		1997 up to now	UNFCCC website
IRR, NPV cost benefit ratio, or unit cost of service	Indicators of investment analysis	Most recent date	Calculation based on local data

Data / parameter	Description	Vintage	Data sources and geographical scale
Investment costs	Including land purchase or rental, machinery, equipments, buildings, fences, site and soil preparation, seedling, planting, weeding, pesticides, fertilization, supervision, training, technical consultation, etc. that occur in the establishment period	Most recent date, taking into account market risk	Local statistics, published data and/or survey
Operations and maintenance costs	Including costs of thinning, pruning, harvesting, replanting, fuel, transportation, repairs, fire and disease control, patrolling, administration, etc.	Most recent date, taking into account market risk	Local statistics, published data and/or survey
Transaction costs	Including costs of project preparation, validation, registration, monitoring, etc.	Most recent date	National and international
Revenues	Those from timber, fuelwood, non-wood products, with and without CER revenues, etc.	Most recent date, taking into account market risk	National or local statistics, published data and/or survey
$C_{BSL,t}$	Baseline net GHG removals by sinks for year t		Estimated per stratum per species
$\Delta C_{ij,BSL,t}$	average annual carbon stock change in the absence of the project activity for year t		Estimated per stratum per species
T	1 to length of crediting period		project
$\Delta C_{ij,t}$	average annual carbon stock change for year t		Estimated per stratum per species
$\Delta C_{G,ij,t}$	average annual increase in carbon due to biomass growth of living trees		Estimated per stratum per species
$\Delta C_{L,ij,t}$	average annual decrease in carbon due to biomass loss of living trees		Estimated per stratum per species
$A_i$	area of baseline stratum i		Estimated per stratum per species
$G_{TOTAL,ij,t}$	average annual increment of total dry biomass of living trees		Estimated per stratum per species
$CF_j$	the carbon fraction		IPCC default, national inventory, literature
44/12	ratio of molecular weights of CO <sub>2</sub> and carbon		IPCC default
$G_{w,ij,t}$	average annual aboveground dry biomass increment of living trees		Estimated per stratum per species
$R_j$	root-shoot ratio		IPCC default, national inventory, literatures
$I_{v,ij,t}$	average annual increment in merchantable volume		IPCC default, national inventory, literatures
$D_j$	basic wood density		IPCC default, national inventory, literatures
$BEF_{1,j}$	biomass expansion factor for conversion of annual net increment (including bark) in stem biomass to total aboveground tree biomass increment		IPCC default, national inventory, literatures

Data / parameter	Description	Vintage	Data sources and geographical scale
$C_{2,ij}$	total carbon stock in living biomass of trees at time 2		Estimated per stratum per species
$C_{1,ij}$	total carbon stock in living biomass of trees at time 1		Estimated per stratum per species
T	number of years between times 2 and 1		
$C_{AB,ij}$	carbon stock in aboveground tree biomass		Estimated per stratum per species
$C_{BB,ij}$	carbon stock in belowground tree biomass		Estimated per stratum per species
$V_{ij}$	merchantable volume of trees		Estimated based on national or local yield table or growth curve
$BEF_{2,j}$	biomass expansion factor for conversion of tree stem biomass to aboveground tree biomass		IPCC Guidelines, IPCC GPG-LULUCF, national inventory, local survey, literature.
$N_{ij}$	Number of trees in baseline scenario		Sampling survey
$f_j(DBH, H)$	An allometric equation linking aboveground biomass of living trees ( $\text{kg d.m tree}^{-1}$ ) to diameter at breast height (DBH) and possibly tree height (H)		Local survey, literatures
$\Delta C_{ijk,t}$	changes in carbon stock in carbon pools		Estimated on stratum/substratum and species basis
$\Delta C_{AB,ijk,t}$	Average annual carbon stock changes in aboveground woody biomass		Estimated on stratum/substratum and species basis
$\Delta C_{BB,ijk,t}$	Average annual carbon stock changes in belowground woody biomass		Estimated on stratum/substratum and species basis
$\Delta C_{SOC,ijk,t}$	Average annual carbon stock changes in soil organic matter		Estimated on stratum/substratum and species basis
$C_{AB,t_2,ijk}$	carbon stock in aboveground woody biomass calculated at time $t_2$		Estimated on stratum/substratum and species basis
$C_{AB,t_1,ijk}$	carbon stock in aboveground woody biomass calculated at time $t_1$		Estimated on stratum/substratum and species basis
$C_{BB,t_2,ijk}$	carbon stock in belowground woody biomass calculated at time $t_2$		Estimated on stratum/substratum and species basis
$C_{BB,t_1,ijk}$	carbon stock in belowground biomass calculated at time $t_1$		Estimated on stratum/substratum and species basis
$C_{AB\_tree,ijk}$	carbon stock in aboveground woody biomass of trees		Estimated on stratum/substratum and species basis

Data / parameter	Description	Vintage	Data sources and geographical scale
$C_{AB\_shrub,ijk}$	carbon stock in aboveground biomass of planted shrubs		Estimated on stratum/substratum and species basis
$C_{BB,tree,ijk}$	carbon stock in below-ground biomass of trees		Estimated on stratum/substratum and species basis
$C_{BB\_shrub,ijk}$	carbon stock in below-ground biomass of planted shrubs		Estimated on stratum/substratum and species basis
$T_1$	number of years between time $t_2$ and $t_1$ for biomass		
$BEF_k$	Species specific tree biomass expansion factor for conversion of merchantable or standing stem biomass to aboveground tree biomass	Most updated	GPG-LULUCF, IPCC 2006 Guidelines, national GHG inventory, local survey
$A_{tree,ijk}$	area covered by trees in a stratum / substratum		Estimated per stratum/substratum per species
$D_k$	biomass expansion factor for conversion of biomass of merchantable or standing volume to above-ground biomass	Most updated	GPG-LULUCF, IPCC 2006 Guidelines, national GHG inventory, local survey
$V_{tree\_ijk}$	mean merchantable/standing volume		Estimated per stratum per species based on local or national growth curve, yield table
$CF_k$	carbon fraction	Most updated	GPG-LULUCF, IPCC 2006 Guidelines, national GHG inventory, local survey, ,per species
$R_k$	Root-shoot ratio of trees	Most updated	GPG-LULUCF, IPCC 2006 Guidelines, national GHG inventory, local survey, per species
$A_{shrub,ijk}$	area covered by shrub in a stratum / substratum		Estimated per stratum/substratum per species
$CF_{s,k}$	carbon fraction of shrub species	Most updated	national GHG inventory, literature, IPCC
$R_{s,k}$	root-shoot ratio of shrub species	Most updated	national GHG inventory, literature, IPCC
$f(DB, H, C, N)$	allometric equation linking above-ground biomass ( $d.m\ ha^{-1}$ ) of shrubs to diameter at base (DB), shrub height (H), crown area/diameter (C) and possibly number of stems (N)	Most updated	national GHG inventory, literature, IPCC
$SOC_{For,ijk}$	stable soil organic carbon stock per hectare of plantation	Most updated	IPCC Guidelines, GPG-LULUCF, national inventory, literature, species specific

Data / parameter	Description	Vintage	Data sources and geographical scale
$SOC_{\text{Non-For},ij}$	stable soil organic carbon stock per hectare of lands before planting	Most updated	IPCC Guidelines, GPG-LULUCF, national inventory, literature, species specific
$T_{\text{For},ij}$	duration of transition from $SOC_{\text{Non-For},ij}$ to $SOC_{\text{For},ijk}$	Most updated	IPCC Guidelines, GPG-LULUCF, national inventory, literature, species specific
$C_{\text{ACTUAL},t}$	Actual net greenhouse gas removals by sinks		Calculated
$GHG_{E,t}$	Increase in GHG emissions by the sources within the project boundary as a result of the implementation of an A/R CDM project activity		Calculated
$E_{\text{FossilFuel},t}$	Emissions from burning of fossil fuels		Estimated
$E_{\text{biomassloss},t}$	Decrease in carbon stock in living biomass of existing non-tree vegetation		Estimated
$N_2O_{\text{Direct}-N_{\text{fertiliser},t}}$	The increase in direct $N_2O$ emission as a result of direct nitrogen application within the project boundary		Estimated
$N_2O_{N_{\text{fixing}}}$	the increase in $N_2O$ emission as a result of planting of N-fixing species and cultivation of N-fixing annual crops within the project boundary		Estimated
$CSP_{\text{diesel},t}$	Amount of diesel consumption		estimated
$CSP_{\text{gasoline},t}$	Amount of gasoline consumption		estimated
$EF_{\text{diesel}}$	Emission factor for diesel	Most recent	IPCC Guideline, GPG 2000, national inventory
$EF_{\text{gasoline}}$	Emission factor for gasoline	Most recent	IPCC Guideline, GPG 2000, national inventory
$B_{\text{non-tree},i}$	Average biomass stock on land to be planted, before the start of a proposed A/R CDM project activity		Local survey
$CF_{\text{non-tree}}$	The carbon fraction of dry biomass in non-tree vegetation	Most recent	IPCC Guidelines, GPG-LULUCF, national GHG inventory, local survey
$F_{\text{SN},t}$	Annual amount of synthetic fertilizer nitrogen adjusted for volatilization as $NH_3$ and $NO_x$		Estimated
$F_{\text{ON},t}$	Annual amount of organic fertilizer nitrogen adjusted for volatilization as $NH_3$ and $NO_x$		Estimated
$EF_1$	Emission Factor for emissions from N inputs		GPG 2000, IPCC Guidelines, national inventory
$Frac_{\text{GASF}}$	The fraction that volatilises as $NH_3$ and $NO_x$ for synthetic fertilizers		GPG 2000, IPCC Guidelines, national inventory

Data / parameter	Description	Vintage	Data sources and geographical scale
$Frac_{GASM}$	The fraction that volatilises as $NH_3$ and $NO_x$ for organic fertilizers		GPG 2000, IPCC Guidelines, national inventory
$N_{SN-Fert,t}$	Amount of synthetic fertilizer nitrogen applied		estimated
$N_{SN-Fert,t}$	Amount of organic fertilizer nitrogen applied		estimated
$F_{BN}$	amount of nitrogen fixed by N-fixing intercrops cultivated annually		Estimated
44/28	ratio of molecular weights of $N_2O$ and nitrogen		Global default
$GWP_{N_2O}$	global warming potential for $N_2O$		with a value of 310 for the first commitment period (IPCC)
$F_{BN,t}$	amount of nitrogen fixed by N-fixing intercrops cultivated annually		Estimated
$F_{SBN,t}$	amount of nitrogen fixed by N-fixing shrubs planted		Estimated
$Crop_{BF_k}$	the seed yield of N-fixing crops	Most recent	estimated
$Crop_{RA_k}$	the ratio of dry matter in the aboveground crop biomass (including residue) to the seed yield	Most recent	IPCC guidelines, GPG 2000, national inventory, literatures
$Crop_{NCRBF_k}$	the fraction of crop biomass that is nitrogen	Most recent	IPCC guideline, GPG 2000, national inventory, literatures
$Shrub_{NCRBF_k}$	the fraction of N-fixing shrub biomass that is nitrogen	Most recent	national inventory, literatures
$\Delta B_{AB\_shrub_{ijk}}$	annual stock change of aboveground biomass		Estimated, literatures
$LF_k$	the ratio of leaf biomass in aboveground biomass of N-fixing shrubs	Most updated	Estimated, literatures
$LK_t$	Total GHG emissions caused by transportation		estimated
$LK_{Vehicle,CO_2,t}$	$CO_2$ emissions caused by transportation		estimated
$LK_{FFL,t}$	GHG emissions from the forage-fed livestock		estimated
$FuelConsumption_{ij,t}$	Consumption of fuel type j of vehicle type i		Estimated
$e_{ij}$	Average litres consumed per kilometer traveled for vehicle type i with fuel type j	Most updated	GPG-2000, IPCC Guidelines, national GHG inventory
$k_{ij}$	Kilometers traveled by each of vehicle type i with fuel type j		estimated
$n_{ij}$	Number of vehicles		estimated
$LK_{CH_4_{FFL,Ferm,t}}$	$CH_4$ emissions from enteric fermentation of the forage-fed livestock to be supported by the project		estimated

Data / parameter	Description	Vintage	Data sources and geographical scale
$LK_{CH_4FLL,manure,t}$	CH <sub>4</sub> emissions from manure management excreted by the forage-fed livestock		estimated
$LK_{N_2OFLL,manure,t}$	N <sub>2</sub> O emissions from manure management excreted by the forage-fed livestock		estimated
EF <sub>1</sub>	CH <sub>4</sub> emission factor for the forage-fed livestock to be supported by the project	Most updated	GPG-2000, IPCC Guidelines, national GHG inventory
$Population_t$	Equivalent number of forage-fed livestock to be supported by the project	Most updated	estimated
$Produc_{Forage,t}$	Production of forage by the project		estimated
DBI	Daily biomass intake for the forage-fed livestock to be supported by the project, kg d.m.head <sup>-1</sup> day <sup>-1</sup>		National inventory, literature, IPCC default
$GWP_{CH_4}$	Global warming potential for CH <sub>4</sub>		with a value of 23 for the first commitment period (IPCC default)
EF <sub>2</sub>	Manure management CH <sub>4</sub> Emission factor for the forage-fed livestock supported by the project	Most updated	GPG-2000, IPCC Guidelines, national GHG inventory
$LK_{N_2OFLL,manure,t}$	Direct N <sub>2</sub> O emissions from manure management due to the forage-fed livestock		estimated
$LK_{Inirect\_N_2O,manure}$	Indirect N <sub>2</sub> O emissions from manure management due to the forage-fed livestock		estimated
Nex	Annual average N excretion per head for the forage-fed livestock to be supported by the project	Most updated	GPG-2000, IPCC Guidelines, national GHG inventory
EF <sub>3</sub>	Emission factor for direct N <sub>2</sub> O emissions from manure management for the forage-fed livestock supported by the project.	Most updated	GPG-2000, IPCC Guidelines, national GHG inventory
EF <sub>4</sub>	Emission factor for N <sub>2</sub> O emissions from atmospheric deposition of nitrogen on soils and water surfaces	Most updated	GPG-2000, IPCC Guidelines, national GHG inventory
Frac <sub>Gas</sub>	Fraction of forage-fed livestock manure nitrogen that volatilises as NH <sub>3</sub> and NO <sub>x</sub> under the selected manure management system for forage-fed livestock to be supported by the project	Most updated	GPG-2000, IPCC Guidelines, national GHG inventory

## 12. Other information

By applying the proposed methodology, the baseline will be developed in a transparent way:

- Widely available published data will be use whenever possible.
- Archives of land use/cover data from around 1990, or from the year 50 years before the start of



the project, and from a most recent date before the start of a proposed A/R CDM project activity are set up and made available to the public. This will ensure that the lands to be planted are explicitly eligible for an A/R CDM project activity.

- Specific geographical positions including the coordinates of the polygons that define the boundary of each parcel of lands are recorded, archived and made available to validators and verifiers, allowing a DOE to validate and verify a proposed A/R CDM project activity.
- Land use/cover maps or satellite images / aerial photographs from around 1990, several time points in the last 50 years and most recent dates before the start of a proposed A/R CDM project activity are available, ensuring that the sites to be planted are visibly eligible to be an A/R CDM project activity.
- Supplementary surveys on historic changes of land use, land cover and land tenure will be conducted by means of field investigation, land-owner interviews, as well as collection of other data source, and made available to validators and verifiers.
- Comments of landowners / land users concerning their barriers of land use will be surveyed and made available.
- Stratification of a proposed project area will be conducted using broad, widely understood national or regional soil and ecosystem classifications, and made available.

### **Section III: Monitoring methodology description**

Monitoring of project implementation includes:

- Monitoring of the project boundary;
- Monitoring of forest establishment;
- Monitoring of forest management.

The corresponding methodology procedures are outlined below.

#### **1. Monitoring project boundary and project implementation**

##### **(a) Monitoring the boundary of the proposed A/R CDM project activity**

This is meant to demonstrate that the actual planting area conforms to the planting area outlined in the project plan. The following activities are foreseen:

- Field survey concerning the actual boundary within which afforestation/reforestation activity has occurred, site by site;
- Measuring geographical positions (latitude and longitude of each corner of polygon sites) using GPS;
- Checking whether the actual boundary is consistent with the description in the PDD;
- If the actual boundary falls outside of the designed boundary in PDD, additional information for lands beyond the designed boundary in PDD shall be provided; the eligibility of these lands as a part of the A/R CDM project activity shall be justified; and the projected baseline scenario shall be demonstrated to be applicable to these lands. Otherwise, these lands shall not be accounted as a part of the A/R CDM project activity. Such changes in boundary shall be communicated to the DOE and subject to validation during the project, e.g. during the first verification event;
- The project boundary shall be monitored periodically all through the crediting period, including through remote sensing as applicable. If the forest area changes during the crediting period, for instance, because deforestation occurs on the project area, the specific location and area of the deforested land shall be identified. Similarly, if the planting on certain lands within the project boundary fails these lands shall be documented.

**(b) Monitoring of forest establishment**

To ensure that the planting quality confirms to the practice described in PDD and is well-implemented, the following monitoring activities shall be conducted in the first three years after planting:

- Confirm that site and soil preparations are implemented based on practice documented in PDD. For instance, no site burning during site preparation, record date, location, area, biomass removed and other measures undertaken;
- Planting: date, location, area, tree species, stand models;
- Fertilization: date, location, area, tree species, amount and type of fertilizer applied during planting;
- Survival checking:
  - The initial survival rate of planted trees shall be counted three months after the planting, and re-planting shall be conducted if the survival rate is lower than 90 percent of final planting density;
  - Final checking three years after the planting action and re-planting shall be conducted if the survival rate is lower than 80 percent;
  - The checking of the survival rate may be conducted using permanent sample plots;
- Weeding checking: check and confirm that the weeding practice is implemented as described in the PDD;
- Survey and check that species and planting including intercropping for each stratum and substratum are in line with the PDD, including the intercropping of N-fixing plants (species, location, area, output, etc);
- Checking N-fixing species.

**(c) Monitoring of forest management**

Forest management practices are important drivers of the GHG balance of the project, and thus shall be monitored. Practices to be monitored include:

- Thinning: specific location, area, tree species, thinning intensity, biomass removed;
- Harvesting: harvest date, location, area, tree species, volume of biomass removed;
- Fertilization: tree species, location, amount and type of fertilizer applied, etc.;
- Checking and confirming that harvested lands are re-planted or re-sowed in the year of harvesting or the following planting season if direct planting or seeding is used;
- Checking and ensuring that good conditions exist for natural regeneration if harvested lands are allowed to regenerate naturally;
- Survey the annual forage output;
- Monitoring of disturbances: date, location, area (GPS coordinates and remote sensing, as applicable), tree species, type of disturbance, biomass lost, implemented corrective measures, change in the boundary of strata and stands.

**2. Stratification and sampling for ex-post calculations****(a) Stratification**

Ex-post stratification is necessary before the first monitoring event because there are possible changes in tree/shrubs species arrangement and planting year, intercropping, fertilization, or site preparation in comparison to the PDD. Unexpected disturbances could occur during the crediting period (e.g. due to fire, pests or disease outbreaks), affecting differently different parts of an originally homogeneous

stratum or stand. Forest management (cleaning, planting, thinning, harvesting, coppicing, re-planting, and forage collection) may be implemented at different intensities, dates and spatial locations than originally planned in the PDD. Furthermore, to respond any differences in growth conditions compared to what was expected, the ex-post stratification should also be conducted after the first monitoring event based on variation in carbon stock change for each stratum and substratum to increase monitoring precision or reduce monitoring cost at the same precision level. For example, it may be that within one stratum the estimated changes in carbon stocks point to the existence of two sub-populations. Also, two different strata may be similar enough to allow their merging into one stratum. The number and allocation of permanent monitoring plots should also be ex-post modified based on the ex-post stratification.

The strata built in section II.3 will be used in the monitoring methodology. However, ex-post stratification will be conducted.

- Ex-post stratification before the first monitoring event: following factors will be considered in the post-stratification:
  - ✓ Data from monitoring of actual project boundary, planting species and planting year;
  - ✓ Other data from monitoring of forest establishment and management, e.g., intercropping, fertilization;
- Ex-post stratification after the first monitoring event based on the following elements:
  - ✓ Unexpected disturbance;
  - ✓ Management activities that are different from the PDD description;
  - ✓ Variation in carbon stock change for each stratum and sub-stratum. Strata or sub-strata will be grouped into one stratum or substratum if they have similar carbon stock, carbon stock change and spatial variation. In contrast, stratum or substratum with high spatial variation of carbon stock or carbon stock change should be divided into two or more strata or substrata.

The possible need for ex-post stratification shall be evaluated at each monitoring event and changes in the strata should be reported to the DOE for verification. Monitoring of strata and stand boundaries shall be one preferably using a Geographical Information System (GIS) which allows for integrating data from different sources (including GPS coordinates and Remote Sensing data). The monitoring of strata and stand boundaries is critical for a transparent and verifiable monitoring of the variable area of stratum/substrata and species, which is of outmost importance for an accurate and precise calculation of net anthropogenic GHG removals by sinks.

## **(b) Sampling**

Permanent sampling plots will be used for sampling over time to measure and monitor changes in carbon stocks of above- and below ground biomass. Permanent sample plots are generally regarded as statistically efficient in estimating changes in forest carbon stocks because there is typically a high covariance between observations at successive sampling events. However, it should be ensured that the plots are treated in the same way as other lands within the project boundary, e.g., during site and soil preparation, weeding, fertilization, irrigation, thinning, etc., and should not be destroyed over the monitoring interval. Ideally, staff involved in management activities should not be aware of the location of monitoring plots. Where local markers are used, these should not be visible.

### ***b.1 Determining sample size***

The number of plots depends on species variation, accuracy and monitoring interval. Neyman's criterion of fixed levels of accuracy and costs in sampling, through establishing permanent monitoring plots of an A/R CDM project activity, is the most adequate statistical tool for determining the necessary number of permanent sampling plots for monitoring the project activity. The total sum of samples (n) is estimated as per a criterion of Neyman of fixed levels of accuracy and costs, according

to Wenger (1984)<sup>26</sup>:

$$n = \left( \frac{t}{E} \right)^2 \left[ \sum_{h=1}^L W_h \cdot s_h \cdot \sqrt{C_h} \right] \cdot \left[ \sum_{h=1}^L W_h \cdot s_h / \sqrt{C_h} \right] \quad (\text{M.1})$$

$$n_h = n \cdot \frac{W_h \cdot s_h / \sqrt{C_h}}{\sum_{h=1}^L W_h \cdot s_h / \sqrt{C_h}} \quad (\text{M.2})$$

where:

$L$	total number of strata
$t_\alpha$	t value for a confidence level (95%)
$E$	allowable error ( $\pm 10\%$ of the mean)
$s_h$	standard deviation of stratum h
$n_h$	number of samples per stratum that is allocated proportional to $W_h \cdot s_h / \sqrt{C_h}$ .
$W_h$	$N_h/N$
$N$	number of total sample units (all stratum), $N = \sum N_h$
$N_h$	number of sample units for stratum h, calculated by dividing the area of stratum h by area of each plot
$C_h$	cost to select a plot of the stratum h

The standard deviation of each stratum ( $s_h$ ) can be determined through ex ante estimates of variance of carbon stock in both biomass and soil organic carbon from section II.7. The t value for 95% confidence is approximately equal to 2 when the number of sample plot is over 30. As the first step, use 2 as the t value and if the resulting n is less than 30, use the new n to get a new t value and conduct recalculation. This process can be repeated until the calculated n is stabilized. The allowable error is a value on a per-plot basis and can be estimated as  $\pm 10\%$  of the expected mean carbon stock in biomass at the end of a rotation and in soil per plot, which can be estimated as part of the ex-ante estimation of the actual net GHG removals by sinks described in the baseline methodology.

It is good practice to reasonably modify the sample size after the first monitoring event based on the actual variation of the carbon stock changes determined from taking the n samples.

### ***b.2 Locating sampling plots among strata/substrata***

To avoid subjective choice of plot locations (plot centres, plot reference points, movement of plot centres to more “convenient” positions), the permanent sample plots shall be located systematically with a random start, which is considered good practice in GPG-LULUCF. This can be accomplished with the help of a GPS in the field. The geographical position (GPS coordinate), administrative location, stratum and sub-stratum series number of each plots shall be recorded and archived. The size of plots depends on the density of trees, in general between 100 m<sup>2</sup> for dense stands and 1000 m<sup>2</sup> for open stands.

Also, it is to be ensured that the sampling plots are distributed as evenly as possible. For example, if one stratum consists of three geographically separated sites, then it is proposed to

- divide the total stratum area by the number of plots, resulting in the average area per plot;
- divide the area of each site by this average area per plot, and assign the integer part of the result to this site. e.g., if the division results in 6.3 plots, then 6 plots are assigned to this site, and 0.3 plots are carried over to the next site, and so on.

<sup>26</sup> Wenger, K.F. (ed). 1984. Forestry handbook (2nd edition). New York: John Wiley and Sons.

***b.3 Monitoring frequency***

Monitoring interval depends on the variability in carbon stocks and the rate of carbon accumulation, i.e., the growth rate of trees as of living biomass. Although the verification and certification shall be carried out every five years after the first verification until the end of the crediting period<sup>27</sup>, monitoring interval may be less than five years. However, to reduce the monitoring cost, the monitoring intervals shall coincide with verification time, i.e., five years of interval. Logically, one monitoring and verification event will take place close to the end of the first commitment period, e.g. in the second half of the year 2012. Project participants shall determine the first monitoring time, taking into account:

- The growth rate of trees and the financial needs of the project activity: the later the date of the first verification, the higher will be the amount of net anthropogenic GHG removals by sinks but the lower the financial net present value of a CER;
- Harvesting events and rotation length: The time of monitoring and subsequent verification and certification shall not coincide with peaks in carbon stocks<sup>28</sup>.

***b.4 Measuring and estimating carbon stock changes over time***

At each monitoring event, the growth of individual trees on plots shall be measured, and soil organic carbon shall be sampled and measured as well. Pre-existing trees should conservatively and consistently with the baseline methodology not be measured and accounted for. Although non-tree vegetation such as herbaceous plants, grasses, and shrubs can occur, usually with biomass less than 10 percent, there is also non-tree vegetation on arid and hyper arid deserts and the baseline scenario has assumed a zero stock change for this non-tree biomass. Therefore, non-tree vegetation will not be measured and accounted. The omission of non-tree biomass makes the monitoring conservative. Even if the initial site preparation results in a removal of non-tree biomass, there is no risk to over-estimate the removals. The carbon stock changes in living biomass on each plot are then estimated through Biomass Expansion Factors (BEF) method or allometric equations method, and soil carbon stock changes are estimated using Reliable Minimum Estimate (RME).

**3. Calculation of ex post baseline net GHG removals by sinks, if required**

The baseline carbon stock changes do not need to be monitored after the project is established, because the accepted baseline approach 22(a) assumes continuation of existing changes in carbon pools within the project boundary from the time of project validation. However, if the project participants choose a renewable crediting period, relevant data necessary for determining the renewed baseline, including net greenhouse gas removals by sinks during the crediting period, shall be collected and archived to determine whether the baseline approach and baseline scenario are still valid or have to be updated. Reasons for a possible need for updating may include:

- National, local and sectoral policies that may influence land use in the absence of the proposed A/R CDM project activity;
- Technical progresses that may change the baseline approach and baseline scenario;
- Climate conditions and other environmental factors that may change to such a degree as to significantly change the successional and disturbance processes or species composition, resulting in, e.g., improved climate conditions or available seed source would make the natural regeneration possible that is not expected to occur for the current baseline scenario;
- Significant changes of political, social and economic situation, making baseline approach and the projection of baseline scenario unreasonable;

<sup>27</sup> Paragraph 32 of decision 19/CP.9

<sup>28</sup> Paragraph 12 of appendix B in decision 19/CP.9

- Existing barriers that may be removed, for instance:
  - Removal of existing investment barriers: Local farmers (communities) can afford to the high establishment investment in the early stage or have chance to get commercial loans from banks for the reforestation activity;
  - Removal of existing technological barriers: Local farmers (communities) get knowledge and skills for producing high quality seedling, successful tree planting, controlling forest fire, pest and disease, etc.
  - Removal of existing institutional barriers (e.g., well-organized institutional instruments to integrate separate households and address technological and financial barriers);
- Market that may change the alternative land use, e.g., significant price rising of wood and non-woody products would make the arid and hyper arid deserts economically attractive in the absence of the proposed AR CDM project activity;
- Check that the baseline net GHG removals by sinks are not under-estimated before the crediting period can be renewed using control plots.

The carbon stock changes in the baseline scenario can be estimated by measuring carbon stock in the above-ground biomass on control plots respectively, at the initial stage and at the end of the crediting period. The control plots shall be established outside the project boundary and serve as proxy and accurately reflect the development of the arid and hyper arid deserts in the absence of the project activity. Measuring the carbon stock change in above-ground biomass is usually sufficient for the purpose of baseline scenario checking. If the carbon stock in aboveground biomass at the end of the crediting period is statistically significantly higher than the carbon stock at the start of the crediting period, the baseline net GHG removals by sinks shall be re-set.

#### 4. Data to be collected and archived for the estimation of baseline net GHG removals by sinks

**Table 1: Data to be collected in making decision on re-setting of baseline, in case of renewable crediting period**

ID number	Data Variable	Source of data	Data Unit	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
	National, local and sectoral policies that may influence land use in the absence of the proposed AR CDM project activity	various	n.a	collected	Start and end of the crediting period	As complete as possible	
	Natural and anthropogenic factors influencing land use, land cover and natural regeneration	various	n.a	collected	Start and end of the crediting period	As complete as possible	
3.4.01	Stratum ID	Stratification, map	Al-pha numeric		20 years	100%	Stratum identification for baseline scenario checking
3.4.02	carbon stock in aboveground biomass at the end of the crediting period	Calculated based on baseline plot measurement	t CO <sub>2</sub>	c	the end of the crediting period	100% of baseline plots	Calculated based on baseline plot measurement for different strata/sub-strata
3.4.03	carbon stock in aboveground biomass at the start of the crediting period	Calculated based on baseline plot measurement	t CO <sub>2</sub>	c	the start of the crediting period	100% of baseline plots	Calculated based on baseline plot measurement for different strata/sub-strata
3.4.04	baseline carbon stock change in aboveground biomass over the crediting period	Calculated	t CO <sub>2</sub> -e yr <sup>-1</sup>	c	20 years	100%	Calculated from 3.4.02 and 3.4.03

## 5. Calculation of ex post actual net GHG removal by sinks

The actual net greenhouse gas removals by sinks represent the sum of the verifiable changes in carbon stocks in the carbon pools within the project boundary, minus the increase in GHG emissions measured in CO<sub>2</sub> equivalents by the sources that are increased as a result of the implementation of an A/R CDM project activity, while avoiding double counting, within the project boundary, attributable to the A/R CDM project activity. Therefore,

$$\Delta C_{ACTUAL,t} = \sum_i \sum_j \sum_k \Delta C_{ijk,t} - GHG_{E,t} \quad (M.3)$$

where:

$\Delta C_{ACTUAL,t}$	actual net greenhouse gas removals by sinks, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$\Delta C_{ijk,t}$	verifiable changes in carbon stock change in carbon pools for stratum i, sub-stratum j species k, tonnes CO <sub>2</sub> yr <sup>-1</sup> for year t
$GHG_{E,t}$	increase in GHG emissions by the sources within the project boundary as a result of the implementation of an A/R CDM project activity, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year t
$t$	1 to end of crediting period

Note: In the equations of this monitoring methodology, sub-strata represent age class (planting time). In case all trees are planted within a single year, sub-strata are not needed.

### a. Verifiable changes in carbon stocks in the carbon pools

Since carbon stock changes in pools of litter and dead wood are ignored in this methodology, the verifiable changes in carbon stock equal to the carbon stock changes in aboveground and belowground woody biomass and soil organic matter within the project boundary, estimated using the following equations<sup>29</sup>

$$\Delta C_{ijk,t} = (\Delta C_{AB,ijk,t} + \Delta C_{BB,ijk,t} + \Delta C_{SOC,ijk,t}) \cdot 44/12 \quad (M.4)$$

$$\Delta C_{AB,ijk,t} = (C_{AB,m_2,ijk} - C_{AB,m_1,ijk}) / T \quad (M.5)$$

$$\Delta C_{BB,ijk,t} = (C_{BB,m_2,ijk} - C_{BB,m_1,ijk}) / T \quad (M.6)$$

$$\Delta C_{SOC,ijk,t} = (C_{SOC,m_2,ijk} - C_{SOC,m_1,ijk}) / T \quad (M.7)$$

where:

$\Delta C_{ijk,t}$	verifiable changes in carbon stock in living woody biomass for stratum i, sub-stratum j species k, tonnes CO <sub>2</sub> yr <sup>-1</sup> in year t
$\Delta C_{AB,ijk,t}$	changes in carbon stock in aboveground woody biomass for stratum i, sub-stratum j species k, tonnes C yr <sup>-1</sup> in year t
$\Delta C_{BB,ijk,t}$	changes in carbon stock in belowground woody biomass for stratum i, sub-stratum j species k, tonnes C yr <sup>-1</sup> in year t
$\Delta C_{SOC,ijk,t}$	changes in carbon stock in soil organic matter for stratum i, substratum j, species k, tonnes C yr <sup>-1</sup> in year t
$C_{AB,m_2,ijk}$	carbon stock in aboveground woody biomass for stratum i, sub-stratum j species k, calculated at monitoring point m <sub>2</sub> , tonnes C
$C_{AB,m_1,ijk}$	carbon stock in aboveground woody biomass for stratum i, sub-stratum j species k, calculated at monitoring point m <sub>1</sub> , tonnes C

<sup>29</sup> Refers to GPG-LULUCF Equation 3.2.3



$C_{BB,m2,ijk}$	carbon stock in belowground woody biomass for stratum i, sub-stratum j species k, calculated at monitoring point $m_2$ , tonnes C
$C_{BB,m1,ijk}$	carbon stock in belowground woody biomass for stratum i, sub-stratum j species k, calculated at monitoring point $m_1$ , tonnes C
$C_{SOC,m2,ijk}$	carbon stock in soil organic matter for stratum i, substratum j, species k, calculated at time $m_2$ , tonnes C
$C_{SOC,m1,ijk}$	carbon stock in soil organic matter for stratum i, substratum j, species k, calculated at time $m_1$ , tonnes C
44/12	ratio of molecular weights of carbon and CO <sub>2</sub> , dimensionless
$T$	number of years between monitoring point $m_2$ and $m_1$ , which in this methodology is 5 years.

### a.1 Measuring and estimating carbon stock changes in living woody biomass

The total carbon stock in living biomass in trees and planted shrubs for each stratum and sub-stratum in each monitoring point (m) is calculated from the area of each stratum and sub-stratum and mean carbon stock in aboveground woody biomass and belowground woody biomass per unit area, given by:

$$C_{AB,m,ijk} = C_{AB\_tree,m,ijk} + C_{AB\_shrub,m,ijk} \quad (\text{M.8})$$

$$C_{BB,m,ijk} = C_{BB\_tree,m,ijk} + C_{BB\_shrub,m,ijk} \quad (\text{M.9})$$

where

$C_{AB,m,ijk}$	carbon stock in above-ground woody biomass at monitoring point m, tonnes C
$C_{AB\_tree,m,ijk}$	carbon stock in above-ground biomass of trees at monitoring point m, tonnes C
$C_{AB\_shrub,m,ijk}$	carbon stock in above-ground biomass of planted shrubs, tonnes C
$C_{BB,m,ijk}$	carbon stock in below-ground woody biomass at monitoring point m, tonnes C
$C_{BB\_tree,m,ijk}$	carbon stock in below-ground biomass of trees at monitoring point m, tonnes C
$C_{BB\_shrub,m,ijk}$	carbon stock in below-ground biomass of planted shrubs at monitoring point m, tonnes C

#### a.1.1 Planted trees

The total carbon stock in biomass living planted trees for each stratum and sub-stratum in each monitoring point (m) is calculated from the area of each stratum and sub-stratum and mean carbon stock in aboveground biomass and belowground biomass per unit area, given by:

$$C_{AB\_tree,m,ijk} = A_{ijk} \cdot MC_{AB\_tree,m,ijk} \quad (\text{M.10})$$

$$C_{BB\_tree,m,ijk} = A_{ijk} \cdot MC_{BB\_tree,m,ijk} \quad (\text{M.11})$$

where:

$C_{AB\_tree,m,ijk}$	carbon stock in above-ground biomass of trees at monitoring point m, tonnes C
$C_{BB\_tree,m,ijk}$	carbon stock in below-ground biomass of trees at monitoring point m, tonnes C
$A_{ijk}$	area of stratum i, sub-stratum j, species k, hectare (ha)
$MC_{AB\_tree,m,ijk}$	mean carbon stock in aboveground tree biomass per unit area for stratum i, sub-stratum j, tree species k, tonnes C ha <sup>-1</sup>
$MC_{BB\_tree,m,ijk}$	mean carbon stock in belowground tree biomass per unit area for stratum i, sub-stratum j, tree species k, tonnes C ha <sup>-1</sup>

The mean carbon stock in aboveground biomass and belowground biomass of living trees per unit area is estimated based on field measurements on permanent plots. This can be estimated using two

methods, i.e., Biomass Expansion Factors (BEF) method and Allometric Equations method. However, the measurement and ex-post estimation shall not include pre-project trees. Since pre-project trees will not always be obviously distinguishable from those planted as part of the project, such trees shall be marked using permanent marker before or at the same time of planting.

### **BEF Method**

**Step 1:** Measuring the diameter at breast height (DBH, at 1.3 m above ground) and preferably height of all the trees in the permanent sample plots above a minimum DBH. The minimum DBH varies depending on tree species and climate, for instance, the minimum DBH may be as small as 2.5 cm in arid environments where trees grow slowly, whereas it could be up to 10 cm for humid environments where trees grow rapidly (GPG-LULUCF).

**Step 2:** Estimating the volume of the commercial component of trees based on locally derived equations, expressed as volume per unit area (e.g., m<sup>3</sup>/ha). It is also possible to combine step 1 and step 2 if there are field instruments (e.g. relascope) that measure volume of each tree directly.

**Step 3:** Choosing BEF and root-shoot ratio: The BEF and root-shoot ratio vary with local environmental conditions, species and age of trees, the volume of the commercial component of trees. These parameters can be determined by either developing a local regression equation or selecting from national inventory, Annex 3A.1 Table 3A.1.10 of GPG LULUCF, updated values in IPCC 2006 Guidelines for AFOLU, or from published sources. If a significant amount of effort is required to develop local BEFs and root-shoot ratio, involving, for instance, harvest of trees, then it is recommended not to use this method but rather to use the resources to develop local allometric equations as described in the allometric method below (refers to Chapter 4.3 in GPG LULUCF). If that is not possible either, national species specific defaults are for BEF and R can be used. Since both BEF and the root-shoot ratio are age dependent, it is desirable to use age-dependent equations. Stemwood volume can be very small in young stands and BEF can be very large, while for old stands BEF is usually significantly smaller. Therefore using average BEF value may result in significant errors for both young stands and old stands. It is preferable to use allometric equations, if the equations are available, and as a second best solution, to use age-dependent BEFs (but for very young trees, multiplying a small number for stemwood with a large number for the BEF can result in significant error).

**Step 4:** Converting the volume of the commercial component of trees into carbon stock in aboveground biomass and belowground biomass via basic wood density, BEF root-shoot ratio and carbon fraction, given by<sup>30</sup>:

$$C_{AB\_tree} = V \cdot D \cdot BEF \cdot CF \quad (M.12)$$

$$C_{BB\_tree} = MC_{AB\_tree} \cdot R \quad (M.13)$$

where:

$C_{AB\_tree}$	carbon stock in aboveground biomass of each tree on plot, tonnes C tree <sup>-1</sup>
$C_{BB\_tree}$	carbon stock in belowground biomass of each tree on plot, tonnes C tree <sup>-1</sup>
$V$	merchantable volume of trees, m <sup>3</sup> tree <sup>-1</sup>
$D$	volume-weighted average wood density, tonnes d.m.m <sup>-3</sup> merchantable volume
$BEF$	tree biomass expansion factor for conversion of biomass of merchantable tree stem volume to aboveground tree biomass, dimensionless.
$CF$	carbon fraction, tonnes C (tonne d.m.) <sup>-1</sup> , IPCC default value = 0.5.
$R$	Root-shoot ratio, dimensionless

<sup>30</sup> Refers to GPG LULUCF Equation 4.3.1

**Step 5:** Calculating plot level carbon stock in aboveground and belowground biomass of living trees:

$$C_{AB\_tree,m,ijk,p} = \sum_{l=1}^{N_{m,ijk,p}} C_{AB\_tree,m,ijk,p,l} \cdot \frac{10000}{A_p} \quad (\text{M.14})$$

$$C_{BB\_tree,m,ijk,p} = \sum_{l=1}^{N_{m,ijk,p}} C_{BB\_tree,m,ijk,p,l} \cdot \frac{10000}{A_p} \quad (\text{M.15})$$

where:

$C_{AB\_tree,m,ijk,p}$	Carbon stock in aboveground biomass of trees on plot $p$ of stratum $i$ sub-stratum $j$ species $k$ at monitoring point $m$ , tonnes C ha <sup>-1</sup>
$C_{BB\_tree,m,ijk,p}$	Carbon stock in belowground biomass of trees on plot $p$ of stratum $i$ sub-stratum $j$ species $k$ at monitoring point $m$ , tonnes C ha <sup>-1</sup>
$C_{AB\_tree,m,ijk,p,l}$	Carbon stock in aboveground biomass of tree $l$ on plot $p$ of stratum $i$ sub-stratum $j$ species $k$ at monitoring point $m$ , tonnes C tree <sup>-1</sup>
$C_{BB\_tree,m,ijk,p,l}$	Carbon stock in belowground biomass of tree $l$ on plot $p$ of stratum $i$ sub-stratum $j$ species $k$ at monitoring point $m$ , tonnes C tree <sup>-1</sup>
$N_{m,ijk,p}$	Number of trees on plot $p$ of stratum $i$ sub-stratum $j$ species $k$ at monitoring point $m$
$A_p$	Area of plot $p$ , m <sup>2</sup>
$l$	Sequence number of trees on plot $p$
10000	Conversion m <sup>2</sup> to hectare

**Step 6:** calculating mean carbon stock in above and belowground biomass per unit area for each stratum/sub-stratum and tree species:

$$MC_{AB\_tree,m,ijk} = \frac{\sum_{p=1}^{P_{ijk}} C_{AB\_tree,m,ijk,p}}{P_{ijk}} \quad (\text{M.16})$$

$$MC_{BB\_tree,m,ijk} = \frac{\sum_{p=1}^{P_{ijk}} C_{BB\_tree,m,ijk,p}}{P_{ijk}} \quad (\text{M.17})$$

where:

$MC_{AB\_tree,m,ijk}$	mean carbon stock in aboveground tree biomass per unit area for stratum $i$ sub-stratum $j$ tree species $k$ at monitoring point $m$ , tonnes C ha <sup>-1</sup>
$MC_{BB\_tree,m,ijk}$	mean carbon stock in belowground tree biomass per unit area for stratum $i$ sub-stratum $j$ tree species $k$ at monitoring point $m$ , tonnes C ha <sup>-1</sup>
$C_{AB\_tree,m,ijk,p}$	Carbon stock in aboveground biomass of trees on plot $p$ of stratum $i$ sub-stratum $j$ species $k$ at monitoring point $m$ , tonnes C ha <sup>-1</sup>
$C_{BB\_tree,m,ijk,p}$	Carbon stock in belowground biomass of trees on plot $p$ of stratum $i$ sub-stratum $j$ species $k$ at monitoring point $m$ , tonnes C ha <sup>-1</sup>
$P_{ijk}$	Number of plots in stratum $i$ sub-stratum $j$ species $k$

**Allometric method**

**Step 1:** As with the step 1 in BEF method, the diameter and preferably height of all trees above some minimum diameter is measured.

**Step 2:** Choosing or establishing appropriate allometric equations.

$$B_{AB\_tree} = f(DBH, H) \quad (M.18)$$

where:

$B_{AB\_tree}$  aboveground biomass of living trees, tonnes d.m. tree<sup>-1</sup>  
 $f(DBH, H)$  allometric equation linking aboveground tree biomass (tonnes d.m. tree<sup>-1</sup>) to diameter at breast height (DBH) and possibly tree height (H).

The allometric equations are preferably local-derived and species-specific. When allometric equations developed from a biome-wide database, such as those in Annex 4A.2, Tables 4.A.1 and 4.A.2 of GPG LULUCF, or updated in IPCC 2006 Guidelines for AFOLU, are used, it is necessary to verify by destructively harvesting, within the project area but outside the sample plots, a few trees of different sizes and estimate their biomass and then compare against a selected equation. If the biomass estimated from the harvested trees is within about ±10% of that predicted by the equation, then it can be assumed that the selected equation is suitable for the project. If this is not the case, it is recommended to develop local allometric equations for the project use. For this, a sample of trees, representing different size classes, is destructively harvested, and its total biomass is determined. The number of trees to be destructively harvested and measured depends on the range of size classes and number of species—the greater the heterogeneity the more trees are required. If resources permit, the carbon content can be determined in the laboratory. Finally, allometric equations are constructed relating the biomass with values from easily measured variables, such as the DBH and total height (see Chapter 4.3 in GPG LULUCF). Also generic allometric equations can be used, as long as it can be proven that they retain a conservative approach, i.e., they underestimate carbon sequestration.

**Step 3:** Calculating carbon stock in living biomass of each tree on plots:

$$C_{AB\_tree} = B_{AB\_tree} \cdot CF \quad (M.19)$$

$$C_{BB\_tree} = C_{AB\_tree} \cdot R \quad (M.20)$$

where:

$C_{AB\_tree}$  carbon stock in aboveground biomass of each tree on plot, tonnes C tree<sup>-1</sup>  
 $C_{BB\_tree}$  carbon stock in belowground biomass of each tree on plot, tonnes C tree<sup>-1</sup>  
 $CF$  carbon fraction, tonnes C (tonne d.m.)<sup>-1</sup>, IPCC default value = 0.5.  
 $R$  Root-shoot ratio, dimensionless

**Step 4:** Using equation (M.14) and (M.15) to estimate plot level carbon stock, followed by applying equation (M.16) and (M.17) to estimate mean carbon stock in aboveground and belowground biomass of living trees per unit area for stratum i substratum j tree species k at monitoring point m in tonnes C ha<sup>-1</sup>.

**a.1.2 Planted Shrubs**

The limited precipitation usually cannot supply sufficient water for the normal growth of trees in semi-arid areas. In this case, shrubs may be planted in mixture with trees to reduce water consumption. The biomass of planted shrubs can be measured and estimated using allometric method.

**Step 1:** Measuring crown area (diameter), height, diameter at base of shrub and number of stems in the permanent sample plots.

**Step 2:** Choosing or establishing appropriate allometric equations for shrubs.

$$B_{AB\_shrub} = f(DB, H, C, N) \quad (M.21)$$

where

$B_{AB\_shrub}$  above-ground biomass of planted shrub, tonnes d.m. ha<sup>-1</sup>  
 $f(DB, H, C, N)$  an allometric equation linking above-ground biomass (d.m. ha<sup>-1</sup>) of shrubs ( $B_{AB\_shrub}$ ) to diameter at base (DB), shrub height (H), crown area/diameter (C) and number of stems (N).

**Step 3:** Estimating carbon stock in above-ground biomass of shrubs using selected allometric equations applied to the measurements in Step 1.

$$C_{AB\_shrub,m,ijk} = A_{shrub,ijk} \cdot B_{AB\_shrub,m,ijk} \cdot CF_{s,k} \quad (M.22)$$

where

$C_{AB\_shrub,m,ijk}$  carbon stock in above-ground biomass of planted shrubs of stratum i, substratum j, and shrub species k at monitoring point m, tonnes C  
 $A_{shrub,ijk}$  area of stratum i, substratum j, and shrub species k, hectare (ha)  
 $B_{AB\_shrub,m,ijk}$  above-ground biomass of planted shrub of stratum i, substratum j, and shrub species k at monitoring point m, tonnes d.m. ha<sup>-1</sup>  
 $CF_{s,k}$  carbon fraction of shrub species k, dimensionless

**Step 4:** Estimating carbon stock in below-ground biomass and total carbon stock in biomass of planted shrubs.

$$C_{BB\_shrub,m,ijk} = C_{AB\_shrub,m,ijk} \cdot R_{s,k} \quad (M.23)$$

where

$C_{BB\_shrub,m,ijk}$  carbon stock in below-ground biomass of planted shrubs of stratum i, substratum j, and shrub species k at monitoring point m, tonnes C  
 $C_{AB\_shrub,m,ijk}$  carbon stock in above-ground biomass of planted shrubs of stratum i, substratum j, and shrub species k at monitoring point m, tonnes C  
 $R_{s,k}$  root-shoot ratio of shrub species k, dimensionless

## a.2. Measuring and estimating carbon stock changes in soil organic matter

**Step 1:** Collecting the soil samples at 0-10 cm, 10-30 cm and optional 30-50 cm soil depth with a soil corer. Cores shall be taken at random located points for each species strip of trees, shrubs and intercrops in each sample plot. The number of cores for each species strip in each sample plot depends on variance of soil carbon and the volume of cores. For cores with 5 cm in diameter and 10 cm in length, five cores should be the minimum.

**Step 2:** The cores are then fully aggregated and mixed for respective strips of tree, shrub and intercrops to reduce the variability, and sub-samples shall be taken for respective depth and strips. If overall ploughing is applied or the disturbed area of soil surface is over 10% of the total surface area, at least five cores shall be randomly taken respectively for ploughed area and non-ploughed area for strips of tree and shrub, and mixed for ploughed and non-ploughed area, respectively.

**Step 3:** Sub-samples are then moved to laboratory, air dried, sieved through 2 mm sieve, and analyzed for soil organic carbon content.

**Step 4:** Separate cores shall be taken next to each of the carbon analysis cores. Precautions should be taken to avoid compression and disturbance. The samples are oven dried at 105°C and

weighed for bulk density determination. The bulk density equals the oven dry weight of soil in the core divided by the core volume after discounting the volume of coarse fraction of >2 mm..

**Step 5:** Calculating the soil organic carbon stock for each plot by multiplying the carbon concentration (percent mass), bulk density, 1 – (% volume of coarse fragments) and soil depth<sup>31</sup>.

$$C_{SOC,m,ijk,p} = DR \cdot \sum_{l=1}^3 (SOCC_{m,ijkl,d1} \cdot BD_{m,ijkl,d1} \cdot F_{m,ijkl,d1} \cdot Depth_l) + (1 - DR) \cdot \sum_{l=1}^3 (SOCC_{m,ijkl,d2} \cdot BD_{m,ijkl,d2} \cdot F_{m,ijkl,d2} \cdot Depth_l) \quad (M.24)$$

where

$C_{SOC,m,ijk,p}$	carbon stock in soil organic matter for stratum i, substratum j, species k plot p at monitoring point m, tonnes C.ha <sup>-1</sup>
$SOCC_{m,ijkl,d1}$	soil organic carbon content on disturbed area for stratum i, substratum j, species k and soil depth l at monitoring point m, g C (100g soil) <sup>-1</sup>
$SOCC_{m,ijkl,d2}$	soil organic carbon content on non-disturbed area for stratum i, substratum j, species k and soil depth l at monitoring point m, g C (100g soil) <sup>-1</sup>
$BD_{m,ijkl,d1}$	soil bulk density on disturbed area for stratum i, substratum j, species k and soil depth l at monitoring point m, g.cm <sup>-3</sup>
$BD_{m,ijkl,d2}$	soil bulk density on non-disturbed area for stratum i, substratum j, species k and soil depth l at monitoring point m, g.cm <sup>-3</sup>
$Depth_l$	sampling depth for stratum i, substratum j, species k and soil depth l, cm
$F_{m,ijkl,d1}$	1 – (% volume of coarse fragments) on disturbed area for stratum i, substratum j, species k and soil depth l at monitoring point m, to adjust the proportion of volumetric sample occupied by the coarse fragment of > 2mm, dimensionless
$F_{m,ijkl,d2}$	1 – (% volume of coarse fragments) on non-disturbed area for stratum i, substratum j, species k and soil depth l at monitoring point m, to adjust the proportion of volumetric sample occupied by the coarse fragment of > 2mm, dimensionless
$DR$	disturbed ratio of surface land area during site preparation, dimensionless

**Step 6:** Calculating the mean soil organic carbon stock

$$MC_{SOC,m,ijk} = \frac{\sum_{p=1}^{P_{ijk}} C_{SOC,m,ijk,p}}{P_{ijk}} \quad (M.25)$$

where,

$MC_{SOC,m,ijk}$	mean carbon stock in the soil organic matter carbon pool in stratum i sub-stratum j species k at monitoring event m, tonnes C ha <sup>-1</sup> .
$C_{SOC,m,ijk,p}$	carbon stock in soil organic matter for stratum i substratum j species k plot p at monitoring point m, tonnes C.ha <sup>-1</sup>
$P_{ijk}$	Number of plot in stratum i sub-stratum j species k

**Step 7:** Estimating the soil organic carbon stock change using equation (M.24) and Reliable Minimum Estimate (RME) approach<sup>32</sup>. Specifically, change in soil organic carbon can be

<sup>31</sup> Refer to equation 4.3.3 in GPG-LULUCF

<sup>32</sup> Refer to page 4.102-4.103 in GPG-LULUCF

estimated by comparing the mean soil organic carbon accumulation between two monitoring periods using the Reliable Minimum Estimate (RME) (Dawkins, 1957)<sup>33</sup>. Under the RME approach, the monitoring results of the plots are pooled to assess the mean at monitoring interval  $m_2$  and  $m_1$ . The change in soil carbon is calculated by subtracting the maximum estimate of the mean at monitoring time  $m_1$  from the minimum mean estimate at monitoring event  $m_2$ . The resulting difference represents the minimum change in mean soil carbon with 95% confidence between the monitoring interval  $m_2$  and  $m_1$ .

$$C_{SOC,m_2,ijk} = (MC_{SOC,m_2,ijk} - 95\%confidenceLevel) \cdot A_{ijk} \quad (M.26)$$

$$C_{SOC,m_1,ijk} = (MC_{SOC,m_1,ijk} + 95\%confidenceLevel) \cdot A_{ijk} \quad (M.27)$$

where

$C_{SOC,m_2,ijk}$	carbon stock in soil organic matter for stratum i, substratum j, species k, calculated at time $m_2$ , tonnes C
$C_{SOC,m_1,ijk}$	carbon stock in soil organic matter for stratum i, substratum j, species k, calculated at time $m_1$ , tonnes C
$MC_{SOC,m_2,ijk}$	mean carbon stock in soil organic matter per hectare for stratum i, substratum j, species k, calculated at time $m_2$ , tonnes C ha <sup>-1</sup>
$MC_{SOC,m_1,ijk}$	mean carbon stock in soil organic matter per hectare for stratum i, substratum j, species k, calculated at time $m_1$ , tonnes C ha <sup>-1</sup>
$A_{ijk}$	Area of stratum i substratum j species k, ha

The initial soil organic carbon stock shall be sampled and estimated before the start of a proposed A/R CDM project activity, using the steps described above, respectively for each stratum.

#### b. GHG emissions by sources

The increase in GHG emission as a result of the implementation of a proposed A/R CDM project activity within the project boundary can be estimated by:

$$GHG_{E,t} = E_{FuelBurn,t} + E_{Biomassloss,t} + N_2O_{N_{fixing},t} + N_2O_{Direct-fertiliser,t} \quad (M.28)$$

where:

$GHG_{E,t}$	the increase in GHG emission as a result of the implementation of a proposed A/R CDM project activity within the project boundary, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year t
$E_{FossilFuel,t}$	the increase in GHG emission as a result of burning of fossil fuels within the project boundary, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year t
$E_{biomassloss,t}$	decrease in carbon stock in living biomass of existing non-tree vegetation tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year t
$N_2O_{N_{fixing},t}$	increase in N <sub>2</sub> O emission as a result of planting of N-fixing species and cultivation of N-fixing annual crops within the project boundary, tonnes CO <sub>2</sub> -e.yr <sup>-1</sup> in year t
$N_2O_{Direct-N_{fertiliser},t}$	increase in direct N <sub>2</sub> O emission as a result of nitrogenous fertiliser application within the project boundary, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year t

<sup>33</sup> Dawkins, H.C. (1957) Some results of stratified random sampling of tropical high forest. Seventh British Commonwealth Forestry Conference 7 (iii) 1-12.

**b.1 GHG emissions from burning of fossil fuel**

In the context of the afforestation or reforestation, the increase in GHG emission by burning of fossil fuels is most likely resulted from machinery use during site preparation, thinning and logging.

**Step 1:** Monitoring the type and amount of fossil fuels consumed in site preparation or logging.

**Step 2:** Choosing emission factors. There are three possible sources of emission factors:

- National emission factors: These emission factors may be developed by national programmes such as national GHG inventory;
- Regional emission factors;
- IPCC default emission factors, provided that a careful review of the consistency of these factors with the country conditions has been made. IPCC default factors may be used when no other information is available.

**Step 3:** Estimating of GHG emissions resulted from the burning of fossil fuel during site preparation and logging. Although some non-CO<sub>2</sub> GHG (CO, CH<sub>4</sub>, NMVOCs) may be released during combustion process, all the released carbon are accounted as CO<sub>2</sub> emissions based on the Revised 1996 IPCC Guidelines for energy:

$$E_{FossilFuel,t} = (CSP_{diesel,t} \cdot EF_{diesel} + CSP_{gasoline,t} \cdot EF_{gasoline}) \cdot 0.001 \quad (M.29)$$

where:

$E_{FossilFuel,t}$	the increase in GHG emission as a result of burning of fossil fuels within the project boundary, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year t
$CSP_{diesel,t}$	amount of diesel consumption, litre (l) yr <sup>-1</sup> in year t
$CSP_{gasoline,t}$	amount of gasoline consumption, l yr <sup>-1</sup> in year t
$EF_{diesel}$	emission factor for diesel, kg CO <sub>2</sub> l <sup>-1</sup>
$EF_{gasoline}$	emission factor for gasoline, kg CO <sub>2</sub> l <sup>-1</sup>
0.001	conversion kg to tonnes

**b.2 Decrease in carbon stock in living biomass of existing non-tree vegetation**

It is assumed that all existing non-tree vegetation will disappear due to site preparation or competition from planted trees.

**Step 1:** Measuring and estimating the above- and below-ground biomass of existing non-tree vegetation. This task shall be conducted before the start of project activity. The herbaceous plants can be measured by simple harvesting techniques. A small frame (either circular or square), usually encompassing about 0.5-1.0 m<sup>2</sup> or less, is used to aid this task. The material inside the frame is cut to ground level and weighed, and the underground part is also dug and weighed. Well-mixed samples are then collected and oven dried to determine dry-to-wet matter ratios. These ratios are then used to convert the entire sample to oven-dry matter. For shrubs, destructive harvesting techniques can also be used to measure the living biomass. An alternative approach, if the shrubs are large, is to develop local shrub allometric equations based on variables such as crown area and height or diameter at base of plant or some other relevant variable (e.g., number of stems in multi-stemmed shrubs). The equations would then be based on regressions of biomass of the shrub versus some logical combination of the independent variables. The independent variable or variables would then be measured in the sampling plots (Refers to Chapter 4.3 in GPG LULUCF).

**Step 2:** Estimating decrease in carbon stock of existing non-tree vegetation

$$E_{biomassloss,t} = \sum_i A_i \cdot B_{non-tree,i} \cdot CF_{non-tree} \cdot 44/12 \quad \forall t = 1 \quad (M.30)$$

$$E_{biomassloss,t} = 0 \quad \forall t > 1$$



where:

$E_{biomassloss,t}$	decrease in carbon stock in living biomass of existing non-tree vegetation tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year t
$A_i$	area of stratum i, ha
$B_{non-tree,i}$	average biomass stock of non-tree vegetation on land to be planted before the start of a proposed A/R CDM project activity for stratum i, tonnes d.m. ha <sup>-1</sup>
$CF_{non-tree}$	carbon fraction of dry biomass in non-tree vegetation, tonnes C (tonne d.m.) <sup>-1</sup>
44/12	ratio of molecular weights of CO <sub>2</sub> and carbon, dimensionless

### b.3 Direct Nitrous oxide emissions from nitrogen fertilization practices

Only direct emissions from nitrogen fertilization are monitored and estimated in this methodology. The method of 1996 IPCC Guideline, GPG-2000, GPG LULUCF and IPCC 2006 Guideline for AFOLU can be used to estimate the N<sub>2</sub>O emissions.

Step 1: Monitoring and estimating the amount of nitrogen in synthetic and organic fertilizer used within the project boundary;

$$N_{SN-Fert,t} = \sum_k A_k \cdot N_{SN-Fert,k,t} \cdot 0.001 \quad (\text{M.31})$$

$$N_{ON-Fert,t} = \sum_k A_k \cdot N_{ON-Fert,k,t} \cdot 0.001 \quad (\text{M.32})$$

where:

$N_{SN-Fert,t}$	total use of synthetic fertiliser within the project boundary, tonnes N yr <sup>-1</sup> in year t
$N_{ON-Fert,t}$	total use of organic fertiliser within the project boundary, tonnes N yr <sup>-1</sup> in year t
$A_k$	area of tree species k with fertilization, ha
$N_{SN-Fert,k,t}$	use of synthetic fertiliser per unit area for tree species k, kg N ha <sup>-1</sup> yr <sup>-1</sup> in year t
$N_{ON-Fert,k,t}$	use of organic fertiliser per unit area for tree species k, kg N ha <sup>-1</sup> yr <sup>-1</sup> in year t
0.001	conversion kg N to tonnes N

**Step 2:** Choosing the fractions of synthetic and organic fertiliser nitrogen that is emitted as NO<sub>x</sub> and NH<sub>3</sub>, and emission factors. As noted in GPG 2000 and 1996 IPCC Guideline, the default emission factor is 1.25 % of applied N, and this value should be used when country-specific factors are unavailable. Project developer may develop specific emission factors that are more appropriate for their project. Specific good practice guidance on how to derive specific emission factors is given in Box 4.1 of GPG 2000. The default values for the fractions of synthetic and organic fertiliser nitrogen that are emitted as NO<sub>x</sub> and NH<sub>3</sub> are 0.1 and 0.2 respectively in 1996 IPCC Guideline<sup>34</sup>. Note that these factors may be updated in IPCC 2006 Guidelines.

**Step 3:** Calculating direct N<sub>2</sub>O emissions from nitrogen fertilization<sup>35</sup>

$$N_2O_{direct-N_{fertilizer,t}} = [(F_{SN,t} + F_{ON,t}) \cdot EF_1] \cdot 44/28 \cdot GWP_{N_2O} \quad (\text{M.33})$$

$$F_{SN,t} = N_{SN-Fert,t} \cdot (1 - \text{Frac}_{GASF}) \quad (\text{M.34})$$

$$F_{ON,t} = N_{ON-Fert,t} \cdot (1 - \text{Frac}_{GASM}) \quad (\text{M.35})$$

<sup>34</sup> Refers to table 4-17 and table 4-18 in 1996 IPCC Guideline

<sup>35</sup> Refers to Equation 3.2.18 in IPCC GPG-LULUCF, Equation 4.22, Equation 4.23 and Equation 4.3.1 in GPG-2000

where:

$N_2O_{direct-N_{fertilizer},t}$	the direct N <sub>2</sub> O emission as a result of nitrogen application within the project boundary during monitoring interval, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year t
$F_{SN,t}$	amount of synthetic fertilizer nitrogen applied adjusted for volatilization as NH <sub>3</sub> and NO <sub>x</sub> , tonnes N yr <sup>-1</sup>
$F_{ON,t}$	annual amount of organic fertilizer nitrogen applied adjusted for volatilization as NH <sub>3</sub> and NO <sub>x</sub> , tonnes N yr <sup>-1</sup>
$N_{SN-Fert,t}$	amount of synthetic fertilizer nitrogen applied, tonnes N yr <sup>-1</sup>
$N_{ON-Fert,t}$	amount of organic fertilizer nitrogen applied, tonnes N yr <sup>-1</sup>
$EF_{N_{inputs}}$	emission factor for emissions from N inputs, tonnes N <sub>2</sub> O-N (tonnes N input) <sup>-1</sup>
$Frac_{GASF}$	the fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for synthetic fertilizers, tonnes NH <sub>3</sub> -N and NO <sub>x</sub> -N (tonnes N input) <sup>-1</sup>
$Frac_{GASEM}$	the fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for organic fertilizers, dimensionless
44/28	ratio of molecular weights of N <sub>2</sub> O and nitrogen, tonnes NH <sub>3</sub> -N and NO <sub>x</sub> -N (tonnes N input) <sup>-1</sup>
$GWP_{N_2O}$	Global Warming Potential for N <sub>2</sub> O (= 310, valid for the first commitment period)

**b.4 GHG emissions from planting of N-fixing species and intercropping of N-fixing annual crop**<sup>36</sup>

$$N_2O_{N_{fixing},t} = (F_{BN,t} + F_{SBN,t}) \cdot EF_1 \cdot 44/28 \cdot GWP_{N_2O} \quad (M.36)$$

$$F_{BN,t} = \sum_i \sum_j \sum_k (Crop_{BF_k,t} \cdot Crop_{RA_k} \cdot Crop_{NCRBF_k} \cdot A_{ijk}) \quad (M.37)$$

$$F_{SBN,t} = \sum_i \sum_j \sum_k (\Delta B_{AB\_shrub,t,ijk} \cdot LF_k \cdot Shrub_{NCRBF_k} \cdot A_{ijk}) \quad (3M.8)$$

where

$N_2O_{N_{fixing},t}$	the increase in N <sub>2</sub> O emission as a result of planting of N-fixing shrubs and intercropping of N-fixing annual crop within the project boundary, tonnes CO <sub>2</sub> -e.yr <sup>-1</sup> in year t
$F_{BN,t}$	amount of nitrogen fixed by N-fixing intercrops cultivated annually, t N.yr <sup>-1</sup> in year t
$F_{SBN,t}$	amount of nitrogen fixed by N-fixing shrubs planted, t N.yr <sup>-1</sup> in year t
$EF_{N_{inputs}}$	emission factor for emissions from N inputs, tonnes N <sub>2</sub> O-N (tonnes N input) <sup>-1</sup>
$Crop_{BF_k,t}$	the seed yield of N-fixing crops per hectare for crop type k, t d.m. ha <sup>-1</sup> yr <sup>-1</sup> in year t
$Crop_{RA_k}$	the ratio of dry matter in the aboveground biomass (including residue) to the seed yield for crop type k, dimensionless
$Crop_{NCRBF_k}$	the fraction of nitrogen in crop biomass for crop type k, kg N (kg dry matter) <sup>-1</sup>
$A_{ijk}$	area of N-fixing intercrops or shrubs for stratum i substratum j crop type or species k, ha
$\Delta B_{AB\_shrub,t,ijk}$	stock change of aboveground biomass for stratum i substratum j shrub species k, t d.m. ha <sup>-1</sup> yr <sup>-1</sup> in year t
$LF_k$	the ratio of leaf biomass to aboveground biomass of shrubs, dimensionless
$Shrub_{NCRBF_k}$	the fraction of nitrogen in N-fixing shrub biomass for species k, kg N (kg dry matter) <sup>-1</sup>

<sup>36</sup> Refers to Equation 4.20 and Equation 4.25-4.27 in IPCC GPG-2000 for Agriculture.

44/28 ratio of molecular weights of N<sub>2</sub>O and nitrogen, dimensionless  
 $GWP_{N_2O}$  global warming potential for N<sub>2</sub>O (= 310, valid for the first commitment period)

Country or local specific value for  $Crop_{RA_k}$ ,  $Crop_{NCRBF_k}$  and  $Shrub_{NCRBF_k}$  shall be used. If country-specific data are not available, the default values may be chosen from Table 4.16 of GPG-2000 and Table 4-19 of the Reference Manual of the IPCC 1996 Guidelines (0.03 kg N (kg dry matter)<sup>-1</sup>), or updated values in IPCC 2006 Guidelines for AFOLU.

Country-specific emission factor ( $EF_{N\_inputs}$ ) should be used where possible, in order to reflect the specific conditions of a country and the agricultural practices involved. If country-specific emission factor is not available,  $EF_{N\_inputs}$  from other countries with comparable management and climatic conditions are good alternatives. Otherwise, the default emission factor ( $EF_{N\_inputs}$ ) is 1.25 % of input N as noted in GPG-2000, or updated value in IPCC 2006 Guidelines for AFOLU.

## 6. Data to be collected and archived for actual net GHG removals by sinks

**Table 2: Data to be collected or used in order to monitor the verifiable changes in carbon stock in the carbon pools within the project boundary from the proposed A/R CDM project activity, and how this data will be archived:**

ID number	Data Variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.1.01	Stratum ID	Alpha numeric	Stratification map		Before the start of the project	100%	Each stratum has a particular combination of soil type, climate, and possibly tree species
3.6.1.02	Sub-stratum ID	Alpha numeric	Stratification map		Before the start of the project	100%	Each sub-stratum has a particular year to be planted under each stratum
3.6.1.03	Confidence level	%			Before the start of the project	100%	For the purpose of QA/QC and measuring and monitoring precision control
3.6.1.04	Precision level	%			Before the start of the project	100%	For the purpose of QA/QC and measuring and monitoring precision control
3.6.1.05	Standard deviation of each stratum			e	Before the start of the project	100%	Used for estimating numbers of sample plots of each stratum and sub-stratum
3.6.1.06	Number of sample plots			c	Before the start of the project	100%	For each stratum and sub-stratum, calculated from 3.6.1.03-3.6.1.05 using equation (1)-(2)
3.6.1.07	Sample plot ID	Alpha numeric	Project and plot map		Before the start of the project	100%	Numeric series ID will be assigned to each permanent sample plot
3.6.1.08	Plot location		Project and plot map and GPS locating	m	5 years	100%	Using GPS to locate before start of the project and at time of each field measurement
3.6.1.09	Tree species		Project design map		5 years	100%	Arranged in PDD
3.6.1.10	Age of plantation	year	Plot measurement	m	5 years	100% sampling plot	Counted since the planted year
3.6.1.12	Diameter at breast height (DBH)	cm	Plot measurement	m	5 year	100% trees in plots	Measuring at each monitoring time per sampling method

ID number	Data Variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.1.13	Tree height	m	Plot measurement	m	5 year	100% trees in plots	Measuring at each monitoring time per sampling method
3.6.1.14	Merchantable volume of each tree on plots	m <sup>3</sup> tree <sup>-1</sup>	Calculated or plot measurement	c/m	5 year	100% of sampling plots	Calculated from 3.6.1.12 and possibly 3.6.1.13 using local-derived equations, or directly measured by field instrument
3.6.1.15	Wood density	t d.m. m <sup>-3</sup>	Local-derived, national inventory, GPG for LULUCF	e	5 year	100% of sampling plots	Local-derived and species-specific value have the priority
3.6.1.16	Tree biomass expansion factor (BEF)	Dimensionless	Local-derived, national inventory, GPG for LULUCF	e	5 year	100% of sampling plots	Local-derived and species-specific value have the priority
3.6.1.17	Carbon fraction	t C.(t d.m.) <sup>-1</sup>	Local, national, IPCC	e	5 year	100% of sampling plots	Local-derived and species-specific value have the priority
3.6.1.18	Root-shoot ratio	Dimensionless	Local-derived, national inventory, GPG for LULUCF	e	5 year	100% of sampling plots	Local-derived and species-specific value have the priority
3.6.1.19	Carbon stock in aboveground biomass of each tree on plots	t C tree <sup>-1</sup>	Calculated from equation	c	5 year	100% of sampling plots	Calculated from equation (M.12) via 3.6.1.14-3.6.1.17, or from equation (M.18) and (M.19) via 3.6.1.12, 3.6.1.13 and 3.6.1.17
3.6.1.20	Carbon stock in belowground biomass of each tree on plots	t C tree <sup>-1</sup>	Calculated from equation	c	5 year	100% of sampling plots	Calculated from equation (M.13) or equation (M.20) via 3.6.1.18-3.6.1.19

ID number	Data Variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.1.21	Area of plot	m <sup>2</sup>	measurement	m	5 year	100% of sampling plots	Area of permanent sampling plot
3.6.1.22	Number of trees in each sampling plot	Alpha numeric	measurement	m	5 year	100% of sampling plots	Area of permanent sampling plot
3.6.1.23	Carbon stock in aboveground biomass of trees on plot <i>p</i>	tonnes C ha <sup>-1</sup>	Calculated	c	5 year	100% of sampling plots	Calculated using equation (M.14) via 3.6.1.19, 3.6.1.21 and 3.6.1.22
3.6.1.24	Carbon stock in belowground biomass of trees on plot <i>p</i>	tonnes C ha <sup>-1</sup>	Calculated	c	5 year	100% of sampling plots	Calculated using equation (M.15) via 3.6.1.20, 3.6.1.21 and 3.6.1.22
3.6.1.25	Mean Carbon stock in aboveground biomass per unit area per stratum per tree species	t C ha <sup>-1</sup>	Calculated	C	5 year	100% of strata and sub-strata	Calculated using equation (M.16) via 3.6.1.06 and 3.6.1.23
3.6.1.26	Mean Carbon stock in belowground biomass per unit area per stratum per tree species	t C ha <sup>-1</sup>	Calculated	C	5 year	100% of strata and sub-strata	Calculated from 3.6.1.06 and 3.6.1.24
3.6.1.27	Area of stratum, sub-stratum and tree species	ha	Stratification map and data	M	5 year	100% of strata and sub-strata	Actual area of each stratum and sub-stratum
3.6.1.28	Carbon stock in aboveground biomass of stratum per tree species	t C	Calculated from equation (10)	C	5 year	100% of strata and sub-strata	Calculated from equation (M.10) via 3.6.1.27 and 3.6.1.25
3.6.1.29	Carbon stock in belowground biomass of stratum per tree species	t C	Calculated from equation (11)	C	5 year	100% of strata and sub-strata	Calculated from equation (M.11) 3.6.1.26 and 3.6.1.27
3.6.1.30	Crown diameter of planted shrub	m	Plot measurement	M	5 years	100% of strata and substrata plot	Measuring all shrubs at each monitoring event for all strata/substrata
3.6.1.31	height of planted shrub	m	Plot measurement	M	5 years	100% of strata and substrata plot	Measuring all shrubs at each monitoring event for all strata/substrata

ID number	Data Variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.1.32	Diameter at base of planted shrub	cm	Plot measurement	M	5 years	100% of strata and substrata plot	Measuring all shrubs at each monitoring event for all strata/substrata
3.6.1.33	Number of stem for each planted shrub	numeric	Plot measurement	M	5 years	100% of strata and substrata plot	Measuring all shrubs at each monitoring event for all strata/substrata
3.6.1.34	Aboveground biomass of planted shrub per ha per stratum/substratum per species	t d.m.ha <sup>-1</sup>	Calculation	C	5 years	100% of strata and substrata	Calculated using equation (M.21) via 3.6.1.30-3.6.1.33
6.1.1.35	Area of planted shrubs per stratum/substratum per species	ha	Stratification map and data	M	5 years	100% of strata and substrata	Actual area for each shrub species of each stratum and substratum
3.6.1.36	Carbon fraction of each shrub species	t C.(t d.m.) <sup>-1</sup>	Local, national, IPCC	E	Before the first monitoring	100% of shrub species	Local-derived and species-specific value have the priority
3.6.1.37	Carbon stock in aboveground biomass of planted shrub per stratum/substratum per species	t C	Calculation	C	5 years	100% of shrub species	Calculated using equation (M.22) via 3.6.1.34-3.6.1.36
3.6.1.38	Root-shoot ratio of shrubs	Dimensionless	Local-derived, national inventory,	e	Before the first monitoring	100% of species	Local-derived and species-specific value have the priority
3.6.1.39	Carbon stock in belowground biomass of planted shrub per stratum/substratum per species	t C	calculation	c	5 years	100% of shrub species	Calculated using equation (M.23) via 3.6.1.37-3.6.1.38
3.6.1.40	Carbon stock in aboveground biomass per stratum/substratum per species	t C	Calculated	c	5 years	100% of strata and substrata	Calculated using equation (M.8) via 3.6.1.28 and 3.6.1.37

ID number	Data Variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.1.41	Carbon stock in belowground biomass per stratum per species	t C	Calculated	c	5 years	100% of strata and substrata	Calculated using equation (M.9) via 3.6.1.29 and 3.6.1.39
3.6.1.42	Carbon stock change in aboveground biomass of stratum per species	t C yr <sup>-1</sup>	Calculated	c	5 year	100% of strata and sub-strata	Calculated from equation (M.5) via 3.6.1.40
3.6.1.43	Carbon stock change in belowground biomass of stratum per species	t C yr <sup>-1</sup>	Calculated	c	5 year	100% of strata and sub-strata	Calculated from equation (M.6) 3.6.1.41
3.6.1.44	Soil organic carbon content	g C (100 g soil) <sup>-1</sup>	Plot sampling and analyzing	m	10-20	100% of strata and substrata	Measured for under each species of each stratum and substratum
3.6.1.45	Soil bulk density	100 g soil. cm <sup>-3</sup>	Plot sampling and measuring	m	10-20	100% of strata and substrata	Measured under each species of each stratum and substratum
3.6.1.46	Soil layer	cm	Plot measurement	m	10-20	100% of strata and substrata	Measured under each species of each stratum and substratum
3.6.1.47	proportion of volumetric sample occupied by the coarse fragment of > 2mm	dimension less	measurement	m	10-20	100% of strata and substrata	
3.6.1.48	disturbed ratio of surface land area during site preparation	dimension less	measurement	m	10-20	100% of strata and substrata	
3.6.1.49	Plot level soil organic carbon stock per ha, per species for each stratum and substratum	t C. ha <sup>-1</sup>	Calculation	c	10-20	100% of strata and substrata	Calculated using equation (M.24) via 3.6.1.44-3.6.1.48
3.6.1.50	Mean soil organic carbon stock per ha, per species for each stratum and substratum	t C. ha <sup>-1</sup>	Calculation	c	10-20	100% of strata and substrata	Calculated using equation (M.25) via 3.6.1.06-3.6.1.49



ID number	Data Variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.1.51	Soil organic carbon stock, per species for each stratum and substratum	tC	Calculation	c	10-20	100% of strata and substrata	Calculated using equation (M.26) and (M.27) via 3.6.1.50
3.6.1.52	Stock change in soil organic matter, per species per stratum/substratum	t C. yr <sup>-1</sup>	Calculation	c	10-20	100% of strata and substrata	Calculated using equation (M.7) via 3.6.1.51
3.6.1.53	Total carbon stock change	t CO <sub>2</sub> -e yr <sup>-1</sup>	Calculated	c	5 year	100% project area	Summing up carbon stock change using equation (M.4) via 3.6.1.43, 3.6.1.42 and 3.6.1.52 for all strata, substrata and species

**Table 3: Data to be collected or used in order to monitor the GHG emissions by the sources, measured in units of CO<sub>2</sub> equivalent, that are increased as a result of the implementation of the proposed A/R CDM project activity within the project boundary, and how this data will be archived:**

<b>ID number</b>	<b>Data variable</b>	<b>Data unit</b>	<b>Data source</b>	<b>Measured (m) calculated (c) estimated (e)</b>	<b>Recording frequency</b>	<b>Proportion of data monitored</b>	<b>Comment</b>
3.6.2.01	Amount of diesel consumed in machinery use for site prep, thinning or logging	litre	On-site monitoring	m	Annually	100%	Measuring either diesel consumption per unit area for site preparation, or per unit volume logged or thinned
3.6.2.02	Amount of gasoline consumed in machinery use for site prep, thinning or logging	litre	On-site monitoring	m	Annually	100%	Measuring either diesel consumption per unit area for site preparation, or per unit volume logged or thinned
3.6.2.03	Emission factor for diesel	kg/ litre	GPG 2000, IPPCC Guidelines, national inventory	e	At beginning of the project	100%	National inventory value should has priority
3.6.2.04	Emission factor for gasoline	kg/ litre	GPG 2000, IPPCC Guidelines, national inventory	e	At beginning of the project	100%	National inventory value should has priority
3.6.2.05	Emission from fossil fuel use within project boundary	t CO <sub>2</sub> -e yr <sup>-1</sup>	Calculated	e	Annually	100%	Calculating using equation (M.29) via 3.6.2.01-3.6.2.04
3.6.2.06	Amount of synthetic fertilizer N applied per unit area	kg N ha <sup>-1</sup> yr <sup>-1</sup>	Monitoring activity	m	Annually	100%	For different tree species or management intensity
3.6.2.07	Amount of organic fertilizer N applied per unit area	kg N ha <sup>-1</sup> yr <sup>-1</sup>	Monitoring activity	m	Annually	100%	For different tree species or management intensity
3.6.2.08	Area of land with N applied	Ha yr <sup>-1</sup>	Monitoring activity	m	Annually	100%	For different tree species or management intensity

ID number	Data variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.2.09	Amount of synthetic fertilizer N applied	t N yr <sup>-1</sup>	Calculated	c	Annually	100%	Calculated using equation (M.31) via 3.6.2.06 and 3.6.2.08
3.6.2.10	Amount of organic fertilizer N applied	t N yr <sup>-1</sup>	Calculated	c	Annually	100%	Calculated using equation (M.32) via 3.6.2.07 and 3.6.2.08
3.6.2.11	Fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for synthetic fertilizers	Dimensionless	GPG 2000, GPG LULUCF, IPCC Guidelines National inventory	e	Before start of monitoring	100%	IPCC default value (0.1) is used if no more appropriate data
3.6.2.12	Fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for organic fertilizers	Dimensionless	GPG 2000, GPG LULUCF, IPCC Guidelines National inventory	e	Before start of monitoring	100%	IPCC default value (0.2) is used if no more appropriate data
3.6.2.13	Emission factor for emission from N input	N <sub>2</sub> O N-input <sup>-1</sup>	GPG 2000, GPG LULUCF, IPCC Guidelines National inventory	e	Before start of monitoring	100%	IPCC default value (1.25%) is used if no more appropriate data
3.6.2.14	Direct N <sub>2</sub> O emission from nitrogen fertilisation	t CO <sub>2</sub> -e yr <sup>-1</sup>	Calculated	c	Annually	100%	Calculated using equation (M.33) via 3.6.2.09-3.6.2.13
3.6.2.15	Area of stratum	ha	Stratification map and data	m	5 year	100% of strata and sub-strata	Actual area of each stratum and sub-stratum
3.6.2.16	Average biomass stock of non-tree vegetation on land to be planted	tonnes d.m.ha <sup>-1</sup>	Site survey	m	Before planting activities	100% of strata and sub-strata	

ID number	Data variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.2.17	Carbon fraction of dry biomass in non-tree vegetation	tonnes C (tonne d.m.) <sup>-1</sup>	IPCC	e	Before planting activities	Global to local	IPCC default 0.5 can be used if local data unavailable
3.6.2.18	Decrease in carbon stock of existing non-tree vegetation due to A/R project implementation	tCO <sub>2</sub>	Calculating	c	Before planting activities	All strata	Calculating using equation (M.30) via 3.6.2.15-3.6.2.17
3.6.2.19	Seed yield of N-fixing crops per ha per crop type	t d.m. ha <sup>-1</sup> yr <sup>-1</sup>	Monitoring	m	Annually	100%	Recording crop product per unit area for each crop type
3.6.2.20	Ratio of aboveground biomass (including residue) to the seed yield for crop type	dimension less	Local, national and species specific, GPG-2000, IPCC 2006 Guideline	e	Before the first monitoring	100%	Local and national values have priority, IPCC default = 2
3.6.2.21	Nitrogen content in crop biomass	dimension less	Local, national and species specific, GPG-2000, IPCC 2006 Guideline	e	Before the first monitoring	100%	Local and national values have priority
3.6.2.22	Area of N-fixing intercrops or shrubs for each species stratum/substratum	ha	Field monitoring	m	3-5	100%	
3.6.2.23	Nitrogen content in N-fixing shrub biomass	dimension less	Local, national and species specific	e	Before the first monitoring	100%	May be from literature review
3.6.2.24	Amount of nitrogen fixed by N-fixing intercrops cultivated annually	t N yr <sup>-1</sup>	Calculation	c	3-5	100%	Calculated using equation (M.37) via 3.6.2.19-3.6.2.22
3.6.2.25	Amount of nitrogen fixed by N-fixing shrubs	t N yr <sup>-1</sup>	Calculation	c	3-5	100%	Calculated using equation (M.38) via 3.6.2.22, 3.6.2.23

ID number	Data variable	Data unit	Data source	Measured (m) calculated (c) estimated (e)	Recording frequency	Proportion of data monitored	Comment
3.6.2.26	Increase in N <sub>2</sub> O emission due to planting of N-fixing species and intercropping of N-fixing annual crops	t CO <sub>2</sub> -e. yr <sup>-1</sup>	Calculation	c	3-5	100%	Calculated using equation (M.36) via 3.6.2.24 and 2.1.1.25
3.6.2.27	Total increase in GHG emission	t CO <sub>2</sub> -e yr <sup>-1</sup>	Calculated using equation (23)	c	Annually	100%	Calculated using equation (M.28) via 3.6.2.05, 3.6.2.14, 3.6.2.18 and 3.6.2.26

## 7. Leakage

Leakage represents the increase in GHG emissions by sources which occurs outside the boundary of an A/R CDM project activity and which is measurable and attributable to the A/R CDM project activity. The possible leakages under the applicability conditions of the proposed methodology that need to be monitored are:

- CO<sub>2</sub> emissions due to fossil fuel combustion from use of transportation outside the project boundary;
- GHG emissions associated with livestock fed with forage produced by project activities (forage-fed livestock). This includes CH<sub>4</sub> emissions due to enteric fermentation, and CH<sub>4</sub> and N<sub>2</sub>O emissions due to management of manure excreted by the forage-fed livestock.

$$LK_t = LK_{Vehicle,CO_2,t} + LK_{FFL,t} \quad (M.39)$$

where:

$LK_t$	Leakage due to the increase in GHG emissions by sources outside the project boundary and attributable to the A/R CDM project activity, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year $t$
$LK_{Vehicle,CO_2,t}$	CO <sub>2</sub> emissions due to fossil fuel combustion from vehicles, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year $t$
$LK_{FFL,t}$	GHG emissions from the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year $t$

### a. Monitoring of Leakage due to fossil fuel consumption

In the context of A/R activities, fossil fuel combustion from vehicle use due to the transportation of seedlings, labour, staff, and harvest products to or from project sites, as a result of the proposed A/R CDM project activity, can be monitored and estimated using the IPCC's bottom-up approach.

**Step 1:** Collect the travelled distance of different types of vehicles using different fuel types.

**Step 2:** Determine emission factors for different types of vehicles using different fuel types. Country-specific emission factors shall be used if available. Default emission factors provided in the revised IPCC 1996 Guidelines and updated in the GPG 2000 and IPCC 2006 Guidelines may be used if there are no locally available data.

**Step 3:** Estimate the CO<sub>2</sub> emissions using the bottom-up approach described in GPG 2000 for the energy sector<sup>37</sup>.

$$LK_{vehicle,t} = \sum_i \sum_j (EF_{ij} \cdot FuelConsumption_{ij,t}) \cdot 0.001 \quad (M.40)$$

$$FuelConsumption_{ij,t} = n_{ij,t} \cdot k_{ij,t} \cdot e_{ij,t} \quad (M.41)$$

where:

$LK_{Vehicle,CO_2,t}$	CO <sub>2</sub> emissions due to fossil fuel combustion from vehicles, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> in year $t$
$i$	Vehicle type
$j$	Fuel type
$EF_{ij}$	Emission factor for vehicle type $i$ with fuel type $j$ , kg CO <sub>2</sub> -e l <sup>-1</sup>
$FuelConsumption_{ij}$	Consumption of fuel type $j$ of vehicle type $i$ , litre yr <sup>-1</sup> in year $t$
$n_{ij}$	Number of vehicle type $i$ used, yr <sup>-1</sup> in year $t$

<sup>37</sup> Refer to Equation 2.5 and Equation 2.6 in IPCC GPG 2000 for energy sector

$k_{ij}$	Kilometres travelled annually by each of vehicle type $i$ with fuel type $j$ , $\text{km yr}^{-1}$ in year $t$
$e_{ij}$	Average consumption of fuel per kilometre travelled for vehicle type $i$ with fuel type $j$ , $\text{litre km}^{-1}$ in year $t$ .

### b. Monitoring of leakage due to forage-fed livestock

To monitor and calculate the leakage of GHG emissions from forage-fed livestock, the following three GHG emissions by sources shall be monitored:

- $\text{CH}_4$  emissions from enteric fermentation by forage-fed livestock;
- $\text{CH}_4$  emissions from manure management for forage-fed livestock;
- $\text{N}_2\text{O}$  emissions from manure management for forage-fed livestock.

$$LK_{FFL,t} = LK_{CH_4_{FFL,Ferm,t}} + LK_{CH_4_{FFL,manure,t}} + LK_{N_2O_{FFL,manure,t}} \quad (\text{M.42})$$

where:

$LK_{FFL,t}$	GHG emissions from forage-fed livestock, tonnes $\text{CO}_2\text{-e yr}^{-1}$ for year $t$
$LK_{CH_4_{FFL,Ferm,t}}$	$\text{CH}_4$ emissions from enteric fermentation by forage-fed livestock, tonnes $\text{CO}_2\text{-e yr}^{-1}$ for year $t$
$LK_{CH_4_{FFL,manure,t}}$	$\text{CH}_4$ emissions from manure management for forage-fed livestock, tonnes $\text{CO}_2\text{-e yr}^{-1}$ for year $t$
$LK_{N_2O_{FFL,manure,t}}$	$\text{N}_2\text{O}$ emissions from manure management for forage-fed livestock, tonnes $\text{CO}_2\text{-e yr}^{-1}$ for year $t$

The leakage due to forage-fed livestock can be monitored and estimated as per IPCC GPG 2000 for agriculture, and the IPCC 2006 Guidelines for AFOLU.

#### Step 1: Establish the *ex post* forage-fed livestock group

As specified as an applicability condition, all forage produced by the project shall have a similar nutritional value and digestibility, and will support only a single livestock group with a single manure management system. If these conditions are not met, this methodology can not be used.

During *ex ante* estimates of leakage emissions, the livestock group and manure management system may have been identified according to knowledge of intended project activities, forage types, and local farming practices. Alternatively, the livestock group may have been selected *ex ante* by household survey, as the group that is fed the largest amount of forage most similar to that to be produced by the project, together with the associated manure management system for that livestock group. Regardless, no matter how the *ex ante* livestock group and manure management system were selected, an *ex post* survey of households using forage produced by the project shall be completed once every crediting period to confirm (or change as appropriate) the forage-fed livestock group and manure management system used for emissions estimates. Data shall be obtained by survey of a random sample of all households receiving forage from the project—at least 30 households or 10% of households, whichever is greater, should be sampled and data presented to substantiate selection of the forage fed livestock group and manure management system used. Characteristics of the forage-fed livestock group that will help select appropriate enteric  $\text{CH}_4$  emission factors should also be determined/checked using data obtained during the *ex post* household survey—including, for example, mean weight, growth rate, and milk production.

**Step 2:** Collect data on forage production. As the forest management and other human intervention within each stratum is homogeneous (in case it is not, separate strata shall be made as per Section II.3.a), the forage production can be estimated as the forage output of each stratum per hectare, multiplied by the stratum area. As the forage production may fluctuate over the year, it is recommended to calculate the average output.

$$\text{Produc}_{\text{Forage},t} = \sum_i \sum_k \text{Produc}_{\text{Forage},ik,t} \cdot A_{ik} \quad (\text{M.43})$$

where

$\text{Produc}_{\text{Forage},t}$	Output of forage by the project, tonnes d.m. yr <sup>-1</sup> in year t
$\text{Produc}_{\text{Forage},ik,t}$	Output of forage by the project per hectare for stratum i, forage species k, tonnes d.m. ha <sup>-1</sup> yr <sup>-1</sup> in year t
$A_{ik}$	Area of stratum i, forage species k, ha

**Step 3:** Assuming all forage produced by the project will be consumed by the forage-fed livestock, which is a conservative approach, the equivalent population size of the livestock group fed with forage produced by the project can be calculated by:

$$\text{Population}_t = \text{Produc}_{\text{Forage},t} / (\text{DBI} \cdot 365) \quad (\text{M.44})$$

where:

$\text{Population}_t$	Equivalent number of forage-fed livestock supported by the project, head for year t
$\text{Produc}_{\text{Forage},t}$	Production of forage by the project, tonnes d.m. yr <sup>-1</sup> in year t
$\text{DBI}$	Daily biomass intake by forage-fed livestock supported by the project, kg d.m.head <sup>-1</sup> day <sup>-1</sup>
365	Number of days per year

It is preferable to determine DBI through household survey. At least 30 households using forage produced by the project to feed the livestock group should be randomly surveyed. The survey shall ensure that all data on daily biomass intake obtained from the survey is for livestock that are fed only on forage produced by the project. If it is not possible to complete a household survey, or forage-fed livestock also consume on a daily basis other types of forage than that produced by the project alone, DBI can also be chosen from Table 4 in section II.8.b.1, or from default data provided by the GPG 2000 and the IPCC 2006 Guidelines for AFOLU.

**Step 4:** Determine the emission factors for each emission source for the forage-fed livestock group, including emission factors for CH<sub>4</sub> emissions from enteric fermentation (EF<sub>1</sub>), emission factors for CH<sub>4</sub> emissions from manure management (EF<sub>2</sub>), emission factors for direct N<sub>2</sub>O emission from manure management (EF<sub>3</sub>), the fraction of total annual N excretion for the livestock group for which manure is managed, and the fraction of managed livestock manure nitrogen that volatilises as NH<sub>3</sub> and NO<sub>x</sub> in the manure management phase. The best estimates of emissions will usually be obtained using country-specific emission factors that have been fully documented in peer-reviewed publications or are from national GHG inventory. If appropriate country-specific emission factors are unavailable, default emission factors presented in Tables 10.10, 10.11, 10.14, 10.15, 10.16, 10.21 and 11.3 of the IPCC 2006 Guidelines for AFOLU can be used. The IPCC default nitrogen excretion rates and default values for volatilization of NH<sub>3</sub> and NO<sub>x</sub> in manure management systems are presented in Tables 10.19 and 10.22, respectively, in the IPCC 2006 Guidelines for AFOLU.

To select default CH<sub>4</sub> emission factors for enteric fermentation it is important to identify the region most applicable to the project area. Scrutinise the tabulations in Annex 10A.1 of the IPCC 2006 Guidelines for AFOLU to ensure that the underlying animal characteristics such as weight, growth rate and milk production used to develop the emission factors are similar to local conditions.



When selecting a default CH<sub>4</sub> emission factor for manure management, be sure to consult the supporting tables in Annex 10A.2 of IPCC 2006 Guidelines for AFOLU, for the distribution of manure management systems and animal waste characteristics used to estimate emissions. Select an emission factor for a region that most closely matches that of the project circumstances.

The default value for EF<sub>4</sub> in equation 10.27 of the IPCC 2006 Guidelines for AFOLU is 0.01 tonnes N<sub>2</sub>O-N (tonnes NH<sub>3</sub>-N and NO<sub>x</sub>-N emitted)<sup>-1</sup>. Country-specific values for EF<sub>4</sub> should be used with great care because of the special complexity of transboundary atmospheric transport: use of default values is recommended. This is because although specific countries may have specific measurements of N deposition and associated N<sub>2</sub>O flux, in many cases the deposited N may not have originated in their country. Similarly, some of the N that volatilises in their country may be transported to and deposited in another country, where different conditions that affect the fraction emitted as N<sub>2</sub>O may prevail.

**Step 5:** Calculate CH<sub>4</sub> emissions from enteric fermentation by, and CH<sub>4</sub> and N<sub>2</sub>O emissions from manure management for, the forage-fed livestock using methods provided by the IPCC GPG 2000 and the IPCC 2006 Guidelines for AFOLU:

$$LK_{CH_4FFL,Ferm,t} = EF_1 \cdot Population_t \cdot 0.001 \cdot GWP_{CH_4} \quad (M.45)$$

$$LK_{CH_4FFL,manure,t} = EF_2 \cdot Population_t \cdot 0.001 \cdot GWP_{CH_4} \quad (M.46)$$

$$LK_{N_2O_{FFL,manure,t}} = LK_{Direct\_N_2O_{FFL,manure,t}} + LK_{Indirect\_N_2O_{FFL,manure,t}} \quad (M.47)$$

$$LK_{Direct\_N_2O_{FFL,manure,t}} = Population_t \cdot Nex \cdot EF_3 \cdot 0.001 \cdot \frac{44}{28} \cdot GWP_{N_2O} \quad (M.48)$$

$$LK_{Indirect\_N_2O_{FFL,manure,t}} = Population_t \cdot Nex \cdot Frac_{Gas} \cdot EF_4 \cdot 0.001 \cdot \frac{44}{28} \cdot GWP_{N_2O} \quad (M.49)$$

where:

$LK_{CH_4FFL,Ferm,t}$	CH <sub>4</sub> emissions from enteric fermentation by the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>
$EF_1$	Enteric CH <sub>4</sub> emission factor for the forage-fed livestock, kg CH <sub>4</sub> head <sup>-1</sup> yr <sup>-1</sup>
$Population_t$	Equivalent number of forage-fed livestock, head for year <i>t</i>
$LK_{CH_4FFL,manure,t}$	CH <sub>4</sub> emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>
$EF_2$	Manure management CH <sub>4</sub> emission factor for the forage-fed livestock, kg CH <sub>4</sub> head <sup>-1</sup> yr <sup>-1</sup>
$LK_{Direct\_N_2O_{FFL,manure,t}}$	Direct N <sub>2</sub> O emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>
$LK_{Indirect\_N_2O_{FFL,manure,t}}$	Indirect N <sub>2</sub> O emissions from manure management for the forage-fed livestock, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year <i>t</i>
$Nex$	Annual average N excretion per head by the forage-fed livestock, kg N head <sup>-1</sup> yr <sup>-1</sup>
$EF_3$	Emission factor for direct N <sub>2</sub> O emission from manure management for the forage-fed livestock, kg N <sub>2</sub> O-N (kg N) <sup>-1</sup>
$EF_4$	Emission factor for N <sub>2</sub> O emissions from atmospheric deposition of nitrogen on soils and water surfaces, kg N <sub>2</sub> O-N (kg NH <sub>3</sub> -N and NO <sub>x</sub> -N emitted) <sup>-1</sup> . Use of the IPCC default factor of 0.01 is recommended
$Frac_{Gas}$	Fraction of managed livestock manure nitrogen that volatilises as NH <sub>3</sub> and NO <sub>x</sub> in the manure management phase for the forage-fed livestock supported by the project, kg NH <sub>3</sub> -N and NO <sub>x</sub> -N (kg N) <sup>-1</sup> .
$GWP_{CH_4}$	Global Warming Potential for CH <sub>4</sub> (= 23 in the 1 <sup>st</sup> C.P.)

$GWP_{N_2O}$	Global Warming Potential for N <sub>2</sub> O (= 310 in the 1 <sup>st</sup> C.P.)
$\frac{44}{28}$	Conversion of (N <sub>2</sub> O-N) emissions to N <sub>2</sub> O emissions, dimensionless
0.001	Conversion of kilograms into tonnes, dimensionless

**Step 6:** Calculate leakage emissions induced by the forage-fed livestock, using equation (M.42) above.

## 8. Data to be collected and archived for leakage

**Table 4: Data and information that will be collected in order to monitor leakage of the proposed A/R CDM project activity:**

ID number	Data Variable	Data unit	Data sources	Measured (m) Calculated (c) estimated (e)	Recording Frequency	Pro-portion of data monitored	Comment
3.8.1.01	Number of each vehicle type used	number	Monitoring of project activity		Annually	100%	Monitoring number of each vehicle type used
3.8.1.02	Emission factors for road transportation	kg CO <sub>2</sub> -e l <sup>-1</sup>	GPG 2000, IPCC Guidelines, national inventory	e	Annually	100%	National or local value has the priority
3.8.1.03	Kilometres travelled by vehicles	km	Monitoring of project activity	m	Annually	100%	Monitoring kilometres for each vehicle type and fuel type used
3.8.1.04	Fuel consumption per km	litre km <sup>-1</sup>	Local data, national data, IPCC	e	5 years	100%	estimated for each vehicle type and fuel type used
3.8.1.05	Fuel consumption for road transportation	litre	Calculated	c	Annually	100%	Calculated using equation (M.41) via 3.8.1.01, 3.8.1.03, 3.8.1.04
3.8.1.06	Leakage due to vehicle use for transportation	t CO <sub>2</sub> -e yr <sup>-1</sup>	Calculated	c	Annually	100%	Calculated using equation (M.40) via 3.8.1.02, 3.8.1.05
3.8.1.07	Area of each stratum and species	ha	Monitoring activity	m/c	5 years	100%	
3.8.1.08	Output of forage per hectare for different stratum and species	tonnes d.m. ha <sup>-1</sup> yr <sup>-1</sup>	monitoring	c	annually	100%	
3.8.1.09	Total output of forage	tonnes d.m. yr <sup>-1</sup>	calculation	c	annually	100%	Calculated using equation (M.43) via 3.8.1.07 and 3.8.1.08

ID number	Data Variable	Data unit	Data sources	Measured (m) Calculated (c) estimated (e)	Recording Frequency	Pro-portion of data monitored	Comment
3.8.1.10	Daily biomass intake for the forage-fed livestock group supported by the project	kg d.m.head <sup>-1</sup> day <sup>-1</sup>	Estimate or survey	e/m	Once	At least 30 household	If household survey is impossible, choosing value from table D
3.8.1.11	Equivalent number of forage-fed livestock supported by the project	head	Calculating	c	Annually	100%	Calculating using equation (M.44) via 3.8.1.09 and 3.8.1.10
3.8.1.12	CH <sub>4</sub> emission factor for enteric fermentation the forage-fed livestock group	kg CH <sub>4</sub> head <sup>-1</sup> yr <sup>-1</sup>	GPG 2000, IPCC Guidelines, national inventory	e	Before the start of project	100%	National or local value has the priority
3.8.1.13	CH <sub>4</sub> emissions from enteric fermentation by the forage-fed livestock	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	Calculating	c	Annually	100%	Calculating using equation (M.45) via 3.8.1.11 -3.8.1.12
3.8.1.14	Manure management CH <sub>4</sub> emission factor for the forage-fed livestock	kg CH <sub>4</sub> head <sup>-1</sup> yr <sup>-1</sup>	GPG 2000, IPCC Guidelines, national inventory	e	Before the start of project	100%	National or local value has the priority
3.8.1.15	CH <sub>4</sub> emissions from manure management for the forage-fed livestock	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	Calculating	c	Annually	100%	Calculating using equation (M.46) via 3.8.1.11 and 3.8.1.14
3.8.1.16	Annual average N excretion per head for the forage-fed livestock	kg N head <sup>-1</sup> yr <sup>-1</sup>	GPG 2000, IPCC Guidelines, national inventory	e	Before the start of project	100%	National or local value has the priority

ID number	Data Variable	Data unit	Data sources	Measured (m) Calculated (c) estimated (e)	Recording Frequency	Pro-portion of data monitored	Comment
3.8.1.17	Emission factor for direct N <sub>2</sub> O emission from manure management for the forage-fed livestock	N <sub>2</sub> O-N (kg N) <sup>-1</sup>	GPG 2000, IPCC Guidelines, national inventory	e	Before the start of project	100%	National or local value has the priority
3.8.1.18	Fraction of managed livestock manure nitrogen that volatilises as NH <sub>3</sub> and NO <sub>x</sub> in the manure management phase for the forage-fed livestock	kg N <sub>2</sub> O-N (kg NH <sub>3</sub> -N and NO <sub>x</sub> -N emitted) <sup>-1</sup>	GPG 2000, IPCC Guidelines, national inventory	e	Before the start of project	100%	National or local value has the priority
3.8.1.19	Emission factor for N <sub>2</sub> O emissions from atmospheric deposition of nitrogen on soil and water surfaces	kg N <sub>2</sub> O-N (kg NH <sub>3</sub> -N and NO <sub>x</sub> -N emitted) <sup>-1</sup>	GPG 2000, IPCC Guidelines	e	Before the start of project	100%	
3.8.1.20	Direct N <sub>2</sub> O emissions from manure management for the forage-fed livestock	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	Calculating	c	Annually	100%	Calculating using equation (M.48)
3.8.1.21	Indirect N <sub>2</sub> O emissions from manure management for the forage-fed livestock	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	Calculating	c	Annually	100%	Calculating using equation (M.49)
3.8.1.22	Direct N <sub>2</sub> O emissions from manure management for the forage-fed livestock	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	Calculating	c	Annually	100%	Calculating using equation (M.47) via 3.8.1.20 and 3.8.1.21
3.8.1.23	Leakage due to the forage-fed livestock	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	Calculating	c	Annually	100%	Calculating using equation (M.42)

ID number	Data Variable	Data unit	Data sources	Measured (m) Calculated (c) estimated (e)	Recording Frequency	Pro-portion of data monitored	Comment
3.8.1.24	Leakage due to the increase in GHG emissions by sources outside the project boundary	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	Calculating	c	Annually	100%	Calculating using equation (M.39) via 3.8.1.06 and 2.1.23

### 9. Ex post net anthropogenic GHG removal by sinks

The net anthropogenic GHG removals by sinks is the actual net GHG removals by sinks minus the baseline net GHG removals by sinks minus leakage, therefore, following general formula can be used to calculate the net anthropogenic GHG removals by sinks of an A/R CDM project activity ( $C_{AR-CDM}$ ), in tonnes CO<sub>2</sub>-e yr<sup>-1</sup>:

$$C_{AR-CDM,t} = \Delta C_{ACTUAL,t} - \Delta C_{BSL,t} - LK_t \quad (M.50)$$

where:

$C_{AR-CDM,t}$	net anthropogenic GHG removals by sinks, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$\Delta C_{ACTUAL,t}$	actual net GHG removals by sinks, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$\Delta C_{BSL,t}$	baseline net GHG removals by sinks, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t
$LK,t$	leakage, tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t

#### Ways of calculating t-CER and l-CER<sup>38</sup>:

t-CERs reflect the *existing stock change at the time of verification* minus project emissions minus leakage (t CO<sub>2</sub>):

$$t-CER(t_v) = C_p(t_v) - C_B(t_v) - \sum_0^{t_v} E_t - \sum_0^{t_v} LK_t \quad (M.51)$$

$$C_p(t_v) - \sum_0^{t_v} E_t = \sum_1^{t_v} \Delta C_{Actual,t} \quad (M.52)$$

$$C_B(t_v) = \sum_1^{t_v} \Delta C_{BSL,t} \quad (M.53)$$

l-CERs reflect the *increment of the stock change* at the time of verification minus project emissions minus leakage compared to the existing stock change at the previous time of verification (t CO<sub>2</sub>):

$$l-CER(t_v) = [C_p(t_v) - C_p(t_v - \kappa)] - [C_B(t_v) - C_B(t_v - \kappa)] - \sum_{t_v-\kappa}^{t_v} E_t - \sum_{t_v-\kappa}^{t_v} LK_t \quad (M.54)$$

$$C_p(t_v) - C_p(t_v - \kappa) - \sum_{t_v-\kappa}^{t_v} E_t = \sum_{t_v-\kappa}^{t_v} \Delta C_{Actual,t} \quad (M.55)$$

$$C_B(t_v) - C_B(t_v - \kappa) = \sum_{t_v-\kappa}^{t_v} \Delta C_{BSL,t} \quad (M.56)$$

where:

$t-CER(t_v)$	t-CERs issued at year of verification $t_v$ (t CO <sub>2</sub> )
$l-CER(t_v)$	l-CERs issued at year of verification $t_v$ (t CO <sub>2</sub> )
$C_p(t_v)$	existing carbon stocks at the year of verification $t_v$ (t CO <sub>2</sub> )
$C_B(t_v)$	estimated carbon stocks of the baseline scenario at year of verification $t_v$ (t CO <sub>2</sub> )
$E(t)$	annual project emissions (t CO <sub>2</sub> )
$LK(t)$	annual leakage (t CO <sub>2</sub> )
$t_v$	year of verification
$\kappa$	time span between two verification occasions (year)

### 10. Uncertainties

Please see section II. 10.

<sup>38</sup> EB22nd meeting report annex 15.

## Section IV: Lists of variables, acronyms and references

## 1. List of variables used in equations:

Table 1: List of variables

Variable	SI Unit	Description
$C_{BSL,t}$	tonnes CO <sub>2</sub> yr <sup>-1</sup>	sum of the changes in carbon stocks in trees for year t
$\Delta C_{ij,BSL,t}$	tonnes CO <sub>2</sub> yr <sup>-1</sup>	average annual carbon stock change for stratum i, species j in the absence of the project activity for year t
$\Delta C_{ij,t}$	tonnes CO <sub>2</sub> yr <sup>-1</sup>	average annual carbon stock change due to biomass growth of pre-project living trees for stratum i species j
$\Delta C_{G,ij,t}$	tonnes CO <sub>2</sub> yr <sup>-1</sup>	average annual increase in carbon due to biomass growth of living trees for stratum i, species j for year t
$\Delta C_{L,ij,t}$	tonnes CO <sub>2</sub> yr <sup>-1</sup>	average annual decrease in carbon due to biomass loss of living trees for stratum i species j
$A_{ij}$	ha	area of stratum i and species j
$G_{TOTAL,ij,t}$	tonnes of dry matter ha <sup>-1</sup> yr <sup>-1</sup>	average annual increment of total dry biomass of living trees for stratum i species j
$CF_j$	tonnes C (tonne d.m.) <sup>-1</sup>	the carbon fraction for species j
$G_{w,ij,t}$	tonnes d.m ha <sup>-1</sup> yr <sup>-1</sup>	average annual aboveground dry biomass increment of living trees for stratum i species j
$R_j$	dimensionless	Root-shoot ratio appropriate to increments for species j
$D_j$	tonnes d.m. m <sup>-3</sup>	basic wood density for species j
$I_{v,ij,t}$	m <sup>3</sup> ha <sup>-1</sup> yr <sup>-1</sup>	average annual increment in merchantable volume for stratum i species j
$BEF_{1,j}$	dimensionless	biomass expansion factor for conversion of annual net increment (including bark) in stem biomass to total aboveground biomass increment for tree species j
$BEF_{2,j}$	dimensionless	biomass expansion factor for conversion of stem biomass to aboveground tree biomass for species j
$C_{2,ij}$	tonnes C	total carbon stock in living biomass of trees for stratum i, species j, calculated at time 2
$C_{1,ij}$	tonnes C	total carbon stock in living biomass of trees for stratum i, species j, calculated at time 1
$V_{ij}$	m <sup>3</sup> ha <sup>-1</sup>	merchantable volume for stratum i, species j
$C_{ACTUAL}$	tonnes CO <sub>2</sub> yr <sup>-1</sup>	actual net greenhouse gas removals by sinks
$T_i$		number of years between times 2 and 1
$C_{AB,ij}$	tonnes C	carbon stock in aboveground tree biomass for stratum i, species j
$C_{BB,ij}$	tonnes C	carbon stock in belowground tree biomass for stratum i, species j
$N_{ij}$	dimensionless	number of trees of species j in stratum i
$f_i(DBH,H)$	kg d.m. tree <sup>-1</sup>	allometric equation linking aboveground biomass of living trees (kg d.m. tree <sup>-1</sup> ) to mean diameter at breast height (DBH) and possibly tree height (H) for species j
$l$	dimensionless	sequence number of tree species j in stratum i
$\Delta C_{ijk,t}$	tonnes CO <sub>2</sub> yr <sup>-1</sup>	changes in carbon stock in carbon pools for stratum i, substratum j, species k



Variable	SI Unit	Description
$\Delta C_{AB,ijk,t}$	tonnes C yr <sup>-1</sup>	changes in carbon stock in aboveground woody biomass for stratum i, substratum j, species k
$\Delta C_{BB,ijk,t}$	tonnes C yr <sup>-1</sup>	changes in carbon stock in belowground woody biomass for stratum i, substratum j, species k
$\Delta C_{SOC,ijk,t}$	tonnes C yr <sup>-1</sup>	changes in carbon stock in soil organic matter for stratum i, substratum j, species k
$C_{AB,t_2,ijk}$	tonnes C	carbon stock in aboveground woody biomass for stratum i substratum j species k, calculated at time t <sub>2</sub>
$C_{AB,t_1,ijk}$	tonnes C	carbon stock in aboveground woody biomass for stratum i substratum j species k, calculated at time t <sub>1</sub>
$C_{BB,t_2,ijk}$	tonnes C	carbon stock in belowground woody biomass for stratum i substratum j species k, calculated at time t <sub>2</sub>
$C_{BB,t_1,ijk}$	tonnes C	carbon stock in belowground woody biomass for stratum i substratum j species k, calculated at time t <sub>1</sub>
$C_{AB\ tree,ijk}$	tonnes C	carbon stock in aboveground biomass of trees
$C_{AB\ shrub,ijk}$	tonnes C	carbon stock in aboveground biomass of planted shrubs
$C_{BB,tree,ijk}$	tonnes C	carbon stock in below-ground biomass of trees, tonnes C
$C_{BB\ shrub,ijk}$	tonnes C	carbon stock in below-ground biomass of planted shrubs
$T_I$	year	number of years between time t <sub>2</sub> and t <sub>1</sub> for biomass, T <sub>1</sub> = t <sub>2</sub> – t <sub>1</sub>
$A_{tree,ijk}$	ha	area covered by trees for stratum i substratum j species k
$V_{tree\ ijk}$	m <sup>3</sup> ha <sup>-1</sup>	mean merchantable/standing volume for stratum i substratum j and species k
$D_k$	tonnes d.m. m <sup>-3</sup> merchantable /standing volume	volume-weighted average wood density for species k
$BEF_k$	dimensionless	biomass expansion factor for conversion of tree biomass of merchantable or standing volume to above-ground biomass
$CF_k$	tonnes C (tonne d.m.) <sup>-1</sup>	carbon fraction, IPCC default value = 0.5
$R_k$	dimensionless	root-shoot ratio
$A_{shrub,ijk}$	ha	area of stratum i substratum j covered by shrub species k
$CF_{s,k}$	dimensionless	carbon fraction of shrub species k
$R_{s,k}$	dimensionless	root-shoot ratio of shrub species k
$f(DB,H,C,N)$	d.m. ha <sup>-1</sup>	an allometric equation linking above-ground biomass of shrubs to one or more of diameter at base (DB), shrub height (H), crown area/diameter (C) and possibly number of stems (N)
$\Delta SOC_{ijk,t}$	tonnes C yr <sup>-1</sup>	average annual carbon stock change in soil organic matter for stratum i substratum j species k
$SOC_{For,ijk}$	tonnes C ha <sup>-1</sup>	stable soil organic carbon stock per hectare of plantation for stratum i substratum j species k
$SOC_{Non-For,ij}$	tonnes C ha <sup>-1</sup>	stable soil organic carbon stock per hectare of lands before planting for stratum i substratum j
$T_{For,ij}$	year	duration of transition from SOC <sub>Non-For,ij</sub> to SOC <sub>For,ijk</sub>
$GHG_{E,t}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	increase in GHG emission as a result of the implementation of the proposed A/R CDM project activity within the project boundary for year t
$E_{FossilFuel,t}$	tonnes CO <sub>2</sub> -e. yr <sup>-1</sup>	increase in GHG emission as a result of burning of fossil fuels within the project boundary

Variable	SI Unit	Description
$E_{biomassloss, t}$	tonnes CO <sub>2</sub>	CO <sub>2</sub> emissions as a result of a decrease in carbon stock in living biomass of existing non-tree vegetation, tonnes CO <sub>2</sub> . This is an initial loss, and therefore accounted once upfront as part of the first monitoring interval, not per year
$N_2O_{N_{fixing}, t}$	tonnes CO <sub>2</sub> -e. yr <sup>-1</sup>	the increase in N <sub>2</sub> O emission as a result of planting of N-fixing shrubs and cultivation of N-fixing annual crops within the project boundary
$N_2O_{Direct-N_{fertiliser}, t}$	tonnes CO <sub>2</sub> -e. yr <sup>-1</sup>	increase in direct N <sub>2</sub> O emission as a result of nitrogen application within the project boundary
$CSP_{diesel, t}$	litre (l) yr <sup>-1</sup>	volume of diesel consumption
$CSP_{gasoline}$	litre (l) yr <sup>-1</sup>	volume of gasoline consumption
$EF_{diesel}$	kg CO <sub>2</sub> l <sup>-1</sup>	emission factor for diesel
$EF_{gasoline}$	kg CO <sub>2</sub> l <sup>-1</sup>	emission factor for gasoline
$B_{non-tree, i}$	tonnes d.m. ha <sup>-1</sup>	average non tree biomass stock on land to be planted, before the start of a proposed A/R CDM project activity
$CF_{non-tree}$	tonnes C (tonne d.m.) <sup>-1</sup>	the carbon fraction of dry biomass in non-tree vegetation
$N_2O_{direct-N_{fertilizer}}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	direct N <sub>2</sub> O emission as a result of nitrogen application within the project boundary
$F_{SN, t}$	tonnes N yr <sup>-1</sup>	mass of synthetic fertilizer nitrogen applied adjusted for volatilization as NH <sub>3</sub> and NO <sub>x</sub>
$F_{ON}$	tonnes N yr <sup>-1</sup>	annual mass of organic fertilizer nitrogen adjusted for volatilization as NH <sub>3</sub> and NO <sub>x</sub>
$N_{SN-Fert, t}$	tonnes N yr <sup>-1</sup> in year t	mass of synthetic fertilizer nitrogen applied
$N_{ON-Fert, t}$	tonnes N yr <sup>-1</sup> in year t	mass of synthetic fertilizer nitrogen applied
$EF_1$	tonnes N <sub>2</sub> O-N (tonnes N input) <sup>-1</sup>	emission factor for emissions from N inputs
$Frac_{GASF}$	tonnes NH <sub>3</sub> -N and NO <sub>x</sub> -N (tonnes N input) <sup>-1</sup>	the fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for synthetic fertilizers
$Frac_{GASM}$	tonnes NH <sub>3</sub> -N and NO <sub>x</sub> -N (tonnes N input) <sup>-1</sup>	the fraction that volatilises as NH <sub>3</sub> and NO <sub>x</sub> for organic fertilizers
$GWP_{N_2O}$	(with a value of 310 for the first commitment period)	global warming potential for N <sub>2</sub> O
$F_{BN, t}$	t N. yr <sup>-1</sup>	amount of nitrogen fixed by N-fixing intercrops cultivated annually
$F_{SBN, t}$	t N. yr <sup>-1</sup>	amount of nitrogen fixed by N-fixing shrubs planted
$EF_{N_{inputs}}$	tonnes N <sub>2</sub> O-N (tonnes N input) <sup>-1</sup>	emission factor for emissions from N inputs
$Crop_{BF_k, t}$	t d.m. ha <sup>-1</sup> yr <sup>-1</sup>	the seed yield of N-fixing intercrops per hectare for crop type k

Variable	SI Unit	Description
$Crop_{RA_k,t}$	dimensionless	the ratio of dry matter in the aboveground biomass (including residue) to the seed yield for crop type k
$Crop_{NCRBF_k}$	dimensionless	the fraction of crop biomass that is nitrogen for crop type k
$\Delta B_{AB\_shrub_{ijk},t}$	t d.m. ha <sup>-1</sup> yr <sup>-1</sup> in year t	annual stock change of aboveground biomass for stratum i, substratum j, shrub species k
$LF_k$	dimensionless	the ratio of leaf biomass in aboveground biomass of N-fixing shrubs
$Shrub_{NCRBF_k}$	dimensionless	the fraction of N-fixing shrub biomass that is nitrogen for species k
$LK_t$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	leakage due to the increase in GHG emissions by sources outside the project boundary and attributable to the A/R CDM project activity
$LK_{Vehicle,CO_2,t}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	GHG emissions due to fossil fuel combustion from vehicles
$LK_{FFL,t}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	GHG emissions from the forage-fed livestock
$EF_{ij}$	kgCO <sub>2</sub> /litre	emission factor for vehicle type i with fuel type j
$FuelConsumption_{ij}$	litres	consumption of fuel type j of vehicle type i
$e_{ij}$	litres/km	Average fuel consumption of vehicle type i with fuel type j
$k_{ij,t}$	km for year t	kilometres travelled by each of vehicle type i with fuel type j
$n_{ij,t}$		number of vehicles for year t
$LK_{CH_4_{FFL,Ferm,t}}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	CH <sub>4</sub> emissions from enteric fermentation by the forage-fed livestock
$LK_{CH_4_{FFL,manure,t}}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	CH <sub>4</sub> emissions from manure management excreted by forage-fed livestock
$LK_{N_2O_{FFL,manure,t}}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	N <sub>2</sub> O emissions from manure management excreted by forage-fed livestock
$EF_1$	kg CH <sub>4</sub> head <sup>-1</sup> yr <sup>-1</sup>	Enteric CH <sub>4</sub> emission factor for the forage-fed livestock
$Population_t$	head	Equivalent number of forage-fed livestock, for year t
$Produc_{Forage,t}$	kg d.m. yr <sup>-1</sup>	Production of forage by the project in year t
$DBI$	kg d.m.head <sup>-1</sup> day <sup>-1</sup>	Daily biomass intake for the forage-fed livestock,
$GWP_{CH_4}$	(with a value of 23 for the first commitment period)	Global warming potential for CH <sub>4</sub>
$EF_2$	kg CH <sub>4</sub> head <sup>-1</sup> yr <sup>-1</sup>	Manure management CH <sub>4</sub> emission factor for the forage-fed livestock
$LK_{Direct\_N_2O_{FFL,manure,t}}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	Direct N <sub>2</sub> O emissions from manure management for the forage-fed livestock, for year t
$LK_{Indirect\_N_2O_{FFL,manure,t}}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	indirect N <sub>2</sub> O emissions from manure management for the forage-fed livestock, for year t
$Nex$	kg N head <sup>-1</sup> yr <sup>-1</sup>	Annual average N excretion per head for the forage-fed livestock

Variable	SI Unit	Description
$EF_3$	kg N <sub>2</sub> O-N (kg N) <sup>-1</sup>	Emission factor for direct N <sub>2</sub> O emission from manure management for the forage-fed livestock
$EF_4$	kg N <sub>2</sub> O-N (kg NH <sub>3</sub> -N and NO <sub>x</sub> -N emitted) <sup>-1</sup>	Emission factor for N <sub>2</sub> O emissions from atmospheric deposition of forage-sourced nitrogen on soils and water surfaces
$Frac_{Gas}$	kg NH <sub>3</sub> -N and NO <sub>x</sub> -N (kg N) <sup>-1</sup>	fraction of managed livestock manure nitrogen that volatilises as NH <sub>3</sub> and NO <sub>x</sub> in the manure management phase for the forage fed livestock
$L$		total number of strata
$t_\alpha$		t value for a confidence level (95%)
$C_{AR-CDM,t}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	net anthropogenic GHG removal by sink for year t
$U_s$	%	percentage uncertainty on the estimate of the mean parameter value
$\mu$		sample mean value of the parameter
$\sigma$		sample standard deviation of the parameter
$U_c$	%	combined percentage uncertainty
$U_{si}$	%	percentage uncertainty on each term of the sum or difference
$C_{si}$		mean value of each term of the sum or difference
$E$		allowable error ( $\pm 10\%$ of the mean)
$s_h$		standard deviation of stratum h
$n_h$		number of samples per stratum that is allocated proportional to $W_h \cdot s_h / \sqrt{C_h}$
$W_h$		$N_h/N$
$N$		number of total sample units (all stratum), $N = \sum N_h$
$N_h$		number of sample units for stratum h, calculated by dividing the area of stratum h by area of each plot
$C_h$		cost to select a plot of the stratum h
$\Delta C_{ACTUAL,t}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup> for year t	actual net greenhouse gas removals by sinks
$\Delta C_{ijk,t}$	tonnes CO <sub>2</sub> yr <sup>-1</sup>	verifiable changes in carbon stock change in carbon pools for stratum i sub-stratum j species k, for year t
$GHG_{E,t}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	increase in GHG emissions by the sources within the project boundary as a result of the implementation of an A/R CDM project activity, in year t
$t$		1 to end of crediting period
$\Delta C_{AB,ijk,t}$	tonnes C yr <sup>-1</sup>	changes in carbon stock in aboveground woody biomass for stratum i sub-stratum j species k, in year t
$\Delta C_{BB,ijk,t}$	tonnes C yr <sup>-1</sup>	changes in carbon stock in belowground woody biomass for stratum i sub-stratum j species k, in year t
$\Delta C_{SOC,ijk,t}$	tonnes C yr <sup>-1</sup>	changes in carbon stock in soil organic matter for stratum i substratum j species k, in year t
$C_{AB,m2,ijk}$	tonnes C	carbon stock in aboveground woody biomass for stratum i sub-stratum j species k, calculated at monitoring point m <sub>2</sub>

Variable	SI Unit	Description
$C_{AB,m1,ijk}$	tonnes C	carbon stock in aboveground woody biomass for stratum i sub-stratum j species k, calculated at monitoring point $m_1$
$C_{BB,m2,ijk}$	tonnes C	carbon stock in belowground woody biomass for stratum i sub-stratum j species k, calculated at monitoring point $m_2$
$C_{BB,m1,ijk}$	tonnes C	carbon stock in belowground woody biomass for stratum i sub-stratum j species k, calculated at monitoring point $m_1$
$C_{SOC,m2,ijk}$	tonnes C	carbon stock in soil organic matter for stratum i substratum j species k, calculated at time $m_2$
$C_{SOC,m1,ijk}$	tonnes C	carbon stock in soil organic matter for stratum i substratum j species k, calculated at time $m_1$
$T$		number of years between monitoring point $m_2$ and $m_1$ , which in this methodology is 5 years
$C_{AB,m,ijk}$	tonnes C	carbon stock in above-ground woody biomass at monitoring point m
$C_{AB\_tree,ijk}$	tonnes C	carbon stock in above-ground biomass of trees at monitoring point m
$C_{AB\_shrub,m,ijk}$	tonnes C	carbon stock in above-ground biomass of planted shrubs
$C_{BB,m,ijk}$	tonnes C	carbon stock in below-ground woody biomass at monitoring point m
$C_{BB,tree,m,ijk}$	tonnes C	carbon stock in below-ground biomass of trees at monitoring point m
$C_{BB\_shrub,m,ijk}$	tonnes C	carbon stock in below-ground biomass of planted shrubs at monitoring point m
$C_{AB\_tree,m,ijk}$	tonnes C	carbon stock in above-ground biomass of trees at monitoring point m, tonnes C
$C_{BB\_tree,m,ijk}$	tonnes C	carbon stock in below-ground biomass of trees at monitoring point m
$MC_{AB\_tree,m,ijk}$	tonnes C ha <sup>-1</sup>	mean carbon stock in aboveground tree biomass per unit area for stratum i, sub-stratum j, tree species k
$MC_{BB\_tree,m,ijk}$	tonnes C ha <sup>-1</sup>	mean carbon stock in belowground tree biomass per unit area for stratum i, sub-stratum j, tree species k
$C_{AB\_tree}$	tonnes C tree <sup>-1</sup>	carbon stock in aboveground biomass of each tree on plot
$C_{BB\_tree}$	tonnes C tree <sup>-1</sup>	carbon stock in belowground biomass of each tree on plot
$V$	m <sup>3</sup> tree <sup>-1</sup>	merchantable volume of each tree on plot
$D$	tonnes d.m. m <sup>-3</sup> merchantable volume	volume-weighted average wood density
$BEF$	dimensionless	tree biomass expansion factor for conversion of biomass of merchantable volume to aboveground tree biomass.
$R$		Root-shoot ratio
$C_{AB\_tree,m,ijk,p}$	tonnes C ha <sup>-1</sup>	Carbon stock in aboveground biomass of trees on plot $p$ of stratum i sub-stratum j species k at monitoring point m
$C_{BB\_tree,m,ijk,p}$	tonnes C ha <sup>-1</sup>	Carbon stock in belowground biomass of trees on plot $p$ of stratum i sub-stratum j species k at monitoring point m
$C_{AB\_tree,m,ijk,p,l}$	tonnes C tree <sup>-1</sup>	Carbon stock in aboveground biomass of tree $l$ on plot $p$ of stratum i sub-stratum j species k at monitoring point m
$C_{BB\_tree,m,ijk,p,l}$	tonnes C tree <sup>-1</sup>	Carbon stock in belowground biomass of tree $l$ on plot $p$ of stratum i sub-stratum j species k at monitoring point m
$N_{m,ijk,p}$		Number of trees on plot $p$ of stratum i sub-stratum j species k at monitoring point m

Variable	SI Unit	Description
$A_p$	$m^2$	Area of plot $p$
$l$		Sequence number of trees on plot $p$
$P_{ijk}$		Number of plots in stratum $i$ sub-stratum $j$ species $k$
$B_{AB\_tree}$	tonnes d.m. $tree^{-1}$	aboveground biomass of living trees
$f(DBH,H)$	d.m. $tree^{-1}$	an allometric equation linking aboveground biomass to diameter at breast height (DBH) and possibly tree height (H).
$A_{shrub,ijk}$	hectare (ha)	area of stratum $i$ substratum $j$ and shrub species $k$
$CF_{s,k}$		carbon fraction of shrub species $k$
$B_{AB\_shrub,m,ijk}$	tonnes d.m. $ha^{-1}$	above-ground biomass of planted shrub of stratum $i$ , substratum $j$ , and shrub species $k$ at monitoring point $m$
$C_{SOC,m,ijk,p}$	tonnes C. $ha^{-1}$	carbon stock in soil organic matter for stratum $i$ substratum $j$ species $k$ plot $p$ at monitoring point $m$
$SOCC_{m,ijkl,d1}$	g C (100g soil) $^{-1}$	soil organic carbon content on disturbed area for stratum $i$ substratum $j$ species $k$ and soil depth $l$ at monitoring point $m$ ,
$SOCC_{m,ijkl,d2}$	g C (100g soil) $^{-1}$	soil organic carbon content on non-disturbed area for stratum $i$ substratum $j$ species $k$ and soil depth $l$ at monitoring point $m$
$BD_{m,ijkl,d1}$	$g\ cm^{-3}$	soil bulk density on disturbed area for stratum $i$ substratum $j$ species $k$ and soil depth $l$ at monitoring point $m$
$BD_{m,ijkl,d2}$	$g\ cm^{-3}$	soil bulk density on non-disturbed area for stratum $i$ substratum $j$ species $k$ and soil depth $l$ at monitoring point $m$
$Depth_l$	cm	sampling depth for stratum $i$ substratum $j$ species $k$ and soil depth $l$
$F_{m,ijkl,d1}$	dimensionless	1 – (% volume of coarse fragments) on disturbed area for stratum $i$ substratum $j$ species $k$ and soil depth $l$ at monitoring point $m$ , to adjust the proportion of volumetric sample occupied by the coarse fragment of $> 2mm$
$F_{m,ijkl,d2}$	dimensionless	1 – (% volume of coarse fragments) on non-disturbed area for stratum $i$ substratum $j$ species $k$ and soil depth $l$ at monitoring point $m$ , to adjust the proportion of volumetric sample occupied by the coarse fragment of $> 2mm$
$DR$	dimensionless	disturbed ratio of surface land area during site preparation
$MC_{SOC,m,ijk}$	tonnes C. $ha^{-1}$	mean carbon stock in the soil organic matter carbon pool in stratum $i$ sub-stratum $j$ species $k$ at monitoring event $m$
$C_{SOC,m1,ijk}$	tonnes C. $ha^{-1}$	carbon stock in soil organic matter for stratum $i$ substratum $j$ species $k$ plot $p$ at monitoring point $m$
$MC_{SOC,m2,ijk}$	tonnes C $ha^{-1}$	mean carbon stock in soil organic matter per hectare for stratum $i$ , substratum $j$ , species $k$ , calculated at time $m_2$
$MC_{SOC,m1,ijk}$	tonnes C $ha^{-1}$	mean carbon stock in soil organic matter per hectare for stratum $i$ , substratum $j$ , species $k$ , calculated at time $m_1$
$N_{SN-Fert,k,t}$	$kg\ N\ ha^{-1}\ yr^{-1}$	use of synthetic fertiliser per unit area for tree species $k$ , in year $t$
$N_{ON-Fert,k,t}$	$kg\ N\ ha^{-1}\ yr^{-1}$	use of organic fertiliser per unit area for tree species $k$ , in year $t$
$GHG_{E,t}$	tonnes $CO_2$ -e $yr^{-1}$	increase in GHG emission as a result of the implementation of a proposed A/R CDM project activity within the project boundary, in year $t$
$E_{FossilFuel,t}$	tonnes $CO_2$ -e $yr^{-1}$	increase in GHG emission as a result of burning of fossil fuels within the project boundary, in year $t$

<b>Variable</b>	<b>SI Unit</b>	<b>Description</b>
$E_{biomassloss,t}$	tonnes CO <sub>2</sub> -e yr <sup>-1</sup>	decrease in carbon stock in living biomass of existing non-tree vegetation in year t
$A_k$	ha	area of tree species k with fertilization

## 2. References:

All references are quoted in footnotes.