

**CDM-MP65-A10**

## Draft Methodological tool

---

# Project and leakage emissions from cultivation of biomass

Version 02.0 - Draft



DRAFT



**United Nations**  
Framework Convention on  
Climate Change

## COVER NOTE

### 1. Procedural background

1. The secretariat workplan for 2014 includes (as part of MAP project 223) simplifications and streamlining of requirements for accounting for leakage emissions from the use of biomass residues/biomass from cultivation. This includes:

- (a) Leakage associated with biomass residues;
- (b) Leakage associated with shift of pre-project activities when cultivating biomass.

### 2. Purpose

2. The purpose of the draft revised tool is to standardize the procedure for identifying and quantifying leakage emissions throughout large scale and small scale methodologies.

### 3. Key issues and proposed solutions

3. The draft revised tool “Leakage and project emissions from biomass” is prepared based on the need for a consistent and viable approach to estimate emissions from biomass residues utilisation.

4. The draft revision does not account for leakage associated with shift of pre-project activities when cultivating biomass, as an analysis results that any deforestation such shift creates will more than eliminate any emission reductions. However, the text was improved to explain how pre-project activities may be accommodated for through their inclusion in the project boundary.

### 4. Impacts

5. Standardization of the procedure for accounting for leakage due to use of biomass residues in various large-scale and small scale methodologies.

### 5. Subsequent work and timelines

6. The Meth Panel, at its 65<sup>th</sup> meeting, agreed on the draft revised tool. After receiving public inputs on the document, the Meth Panel will continue working on the draft revised tool, at its 66<sup>th</sup> meeting, for recommendation to the Board at a future meeting of the Board.

### 6. Recommendations to the Board

7. Not applicable (call for public input).

| <b>TABLE OF CONTENTS</b>   | <b>Page</b> |
|--|-------------|
| <b>1. INTRODUCTION</b> .....   | <b>4</b>    |
| <b>2. SCOPE, APPLICABILITY, AND ENTRY INTO FORCE</b> .....   | <b>4</b>    |
| 2.1. Scope .....   | 4           |
| 2.2. Applicability .....   | 4           |
| 2.3. Entry into force .....  | 5           |
| <b>3. NORMATIVE REFERENCES</b> .....   | <b>5</b>    |
| <b>4. DEFINITIONS</b> .....  | <b>5</b>    |
| <b>5. PROJECT EMISSION FROM CULTIVATION OF BIOMASS</b> .....   | <b>7</b>    |
| 5.1. Emissions resulting from soil management .....  | 7           |
| 5.1.1. Emissions resulting from loss of soil organic carbon.....   | 8           |
| 5.1.2. Emissions resulting from soil fertilization and management.....   | 9           |
| 5.1.3. Emissions resulting from soil amendment .....   | 9           |
| 5.2. Emissions resulting from energy consumption .....   | 10          |
| 5.3. Emissions resulting from clearance or burning of biomass .....  | 10          |
| <b>6. DETERMINATION OF THE ALTERNATIVE SCENARIO OF THE BIOMASS RESIDUES IN ABSENCE OF THE PROJECT ACTIVITY</b> ..... | <b>11</b>   |
| <b>7. LEAKAGE DUE TO BIOMASS RESIDUES</b> .....  | <b>11</b>   |
| <b>8. LEAKAGE DUE TO SHIFT OF PRE-PROJECT ACTIVITIES</b> .....   | <b>14</b>   |
| <b>9. MONITORING METHODOLOGY</b> .....   | <b>15</b>   |
| 9.1. Data and parameters not monitored .....   | 15          |
| 9.2. Data and parameters monitored .....   | 16          |
| <b>APPENDIX 1. DEFAULT VALUES FOR BIOMASS CULTIVATION</b> .....  | <b>22</b>   |
| <b>APPENDIX 2. EXAMPLE IDENTIFICATION OF ALTERNATIVE USES</b> .....  | <b>26</b>   |
| <b>APPENDIX 3. EXPLANATION OF FACTORS USED</b> .....   | <b>27</b>   |

## 1. Introduction

1. This tool:

- (a) Provides steps—a procedure to determine the project emissions resulting from cultivation of biomass in a dedicated plantation;
- (b) Provides methodological guidance to estimate leakage emissions caused by diversion, to the project activity, of biomass from applications not related to the underlying CDM project.

**Table 1. Parameters determined**

| Parameter   | Unit                | Description   |
|-------------|---------------------|---|
| $PE_{BC,y}$ | t CO <sub>2</sub> e | Project emissions resulting from cultivation of biomass in a dedicated plantation, in year y                                |
| $LE_{BC,y}$ | t CO <sub>2</sub> e | Leakage emissions resulting from cultivation of biomass in a dedicated plantation or from utilization of biomass, in year y |

## 2. Scope, applicability, and entry into force

### 2.1. Scope

2. This tool can be used for estimation of project and leakage emissions resulting from cultivation of biomass in a dedicated plantation of a CDM project activity that uses biomass as a source of energy.
3. This tool also includes approaches for identifying and estimating leakage emissions from project activities that utilise biomass residues as input material.
4. Only positive leakage, i.e. increased emissions outside the project boundary are covered by this tool. If the biomass residues utilization results in any reduction of greenhouse gas (GHG) emissions outside the project boundary, they will not be accounted for.

### 2.2. Applicability

5. For project activities which include biomass cultivation:

- (a) The land in which biomass is cultivated:
  - (i) Does not contain wetlands;
  - (ii) Does not contain organic soils;
  - (iii) Does not contain forests;
  - (iv) Is not subjected to flood irrigation.

6. Pre-project activities are not shifted outside the project boundary, to avoid indirect land use changes as a result of the project activity. This condition is met if one of the following two conditions apply:

(b) The plantation area was abandoned land prior to the implementation of the project activity;

(c) The plantation area was used prior to the implementation of the project area but the pre-project land use of the plantation area will be accommodated for, providing at least the same level of service during the project activity, within the land area included in the project boundary. This could be achieved, inter alia, in the following ways:

(v) At least the same number of cattle as prior to the implementation of the project activity will continue being grazed during the project activity within the land area included in project boundary;

(vi) Due to more efficient farming practice, the pre-project crops can be grown on a smaller area, which is included in the land area included in the project boundary, to achieve the same level of annual production of crops, freeing land for the dedicated plantation;

(d) Settlements are not removed from the land area included in the project boundary.

(b) Desalination is not a substantial source of water in the host country.

6. The tool is also applicable if biomass residues are consumed in a CDM project activity. These could be:

(a) Procured by the project proponents; or

(b) The result of an agro-industrial process under the control of the project proponents.

### 2.3. Entry into force

7. Not applicable (call for public input).

## 3. Normative references

8. This tool refers to the following documents:

(a) "Tool to calculate project or leakage CO<sub>2</sub> emissions from fossil fuel combustion";

(b) "Tool to calculate baseline, project and/or leakage emissions from electricity consumption";

## 4. Definitions

9. The definitions contained in the Glossary of CDM terms shall apply.

10. For the purpose of this tool, the following definitions apply:

- (a) **Abandoned land** - land on which no land management (including crops, forestry, agroforestry and grazing) has been practiced for the longer period of:
- (i) Three years;
  - (ii) The length of one crop rotation of the most recent crop grown on the plot.
- (b) **Forest**<sup>1</sup> - in the absence of national definition of forests, the following shall apply: Forest is a minimum area of land of 0.05–1.0 hectares with tree crown cover (or equivalent stocking level) of more than 10–30 per cent with trees with the potential to reach a minimum height of 2–5 metres at maturity in situ. A forest may consist either of closed forest formations where trees of various storeys and undergrowth cover a high portion of the ground or open forest. Young natural stands and all plantations which have yet to reach a crown density of 10–30 per cent or tree height of 2–5 metres are included under forest, as are areas normally forming part of the forest area which are temporarily unstocked as a result of human intervention such as harvesting or natural causes but which are expected to revert to forest;
- (c) **Indirect land use change** - is land-use change that may be induced on land areas not included in the project boundary as a result of shifting of pre-project activities;
- (d) **Organic soil**<sup>2</sup> - soils are organic if they satisfy the requirements 1 and 2, or 1 and 3 below:
- (i) Thickness of 10 cm or more. A horizon less than 20 cm thick must have 12 per cent or more organic carbon when mixed to a depth of 20 cm;
  - (ii) If the soil is never saturated with water for more than a few days, and contains more than 20 per cent (by weight) organic carbon (about 35 per cent organic matter);
  - (iii) If the soil is subject to water saturation episodes and has either:
    - a. At least 12 per cent (by weight) organic carbon (about 20 per cent organic matter) if it has no clay; or
    - b. At least 18 per cent (by weight) organic carbon (about 30 per cent organic matter) if it has 60 per cent or more clay; or
    - c. An intermediate, proportional amount of organic carbon for intermediate amounts of clay;
- (e) **Pre-project activities** - the land use prior to the implementation of the project activity, considering both land use practices and the primary and final products of the practices. This includes, for example, grazing, cultivation of crops, agroforestry, collection of biomass;

---

<sup>1</sup> As defined by the minimum values of tree crown cover, land area, and tree height selected by the host party for the purposes of the CDM (see paragraph 8, annex to decision 5/CMP.1).

<sup>2</sup> As defined in “Annex A: Glossary” of the 2003 IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry.

- (f) **Project region** – area 250km around the project activity;
- (g) **Stratum** - area of land with uniform properties;
- (h) **Wetland**<sup>3</sup> - this category includes land that is covered or saturated by water for all or part of the year (e.g. peatland) and that does not fall into the forest land, cropland, grassland or settlements categories. This category can be subdivided into managed and unmanaged according to national definitions. It includes reservoirs as a managed sub-division and natural rivers and lakes as unmanaged sub-divisions;

## 5. Project emissions from cultivation of biomass

11. Project emissions resulting from cultivation of biomass in a dedicated plantation are estimated as follows:

$$PE_{BC,y} = PE_{SM,y} + PE_{EC,y} + PE_{BB,y} \quad \text{Equation (1)}$$

Where:

- $PE_{BC,y}$  = Emissions resulting from cultivation of biomass in a dedicated plantation, in year  $y$  (t CO<sub>2</sub>e)
- $PE_{SM,y}$  = Emissions resulting from soil management, in year  $y$  (t CO<sub>2</sub>e)
- $PE_{EC,y}$  = Emissions resulting from energy consumption, in year  $y$  (t CO<sub>2</sub>e)
- $PE_{BB,y}$  = Emissions resulting from burning of biomass, in year  $y$  (t CO<sub>2</sub>e)

12. Biomass originating from land areas included in registered A/R project activities may be considered to have no project emissions, provided that the emission reductions from the A/R project activity have been verified and issued for the time period in which the biomass was harvested.

### 5.1. Emissions resulting from soil management

13. Emissions resulting from soil management are estimated as follows:

$$PE_{SM,y} = PE_{SOC,y} + PE_{SF,y} + PE_{SA,y} \quad \text{Equation (2)}$$

Where:

- $PE_{SM,y}$  = Emissions resulting from soil management, in year  $y$  (t CO<sub>2</sub>e)
- $PE_{SOC,y}$  = Emissions resulting from loss of soil organic carbon, in year  $y$  (t CO<sub>2</sub>e)

<sup>3</sup> As defined in “Annex A: Glossary” of the 2003 IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry.

$PE_{SF,y}$  = Emissions resulting from of soil fertilization and management, in year  $y$  (t CO<sub>2</sub>e)

$PE_{SA,y}$  = Emissions resulting from soil amendment (liming), in year  $y$  (t CO<sub>2</sub>e)

### 5.1.1. Emissions resulting from loss of soil organic carbon

14. To estimate emissions resulting from loss of soil organic carbon, the areas of land are stratified according to:

- (a) Climate region and soil types given in Table 1 in appendix 1;
- (b) Land-use and land management activities on croplands given in Tables 2 and 3 in appendix 1; and
- (c) Land-use and land management activities on grasslands given in Table 4 in appendix 1. These apply also to abandoned land.

15. Emissions resulting from loss of soil organic carbon are estimated as follows:

$$PE_{SOC,y} = \max\left(\frac{44}{12} \times \frac{1.156}{T} \times \sum_i \Delta SOC_i, 0\right) \quad \text{Equation (3)}$$

Where:

$PE_{SOC,y}$  = Emissions resulting from loss of soil organic carbon, in year  $y$  (t CO<sub>2</sub>e)

$T$  = Length of the first crediting period of the project in years (7 or 10)

$\Delta SOC_i$  = Loss of soil organic carbon in land stratum  $i$ , (t C)

$\frac{44}{12}$  = Factor for converting units from t C to t CO<sub>2</sub>e; dimensionless

1.156 = Factor to account for soil N<sub>2</sub>O emissions associated with loss of soil organic carbon,<sup>4</sup> dimensionless

$i$  = Strata of areas of land

16. Loss of soil organic carbon in a stratum is estimated as follows:

$$\Delta SOC_i = 1.21 \times A_{SOC,i} \times SOC_{REF,i} \times (f_{LUB,i} \times f_{MGB,i} \times f_{INB,i} - f_{LUP,i} \times f_{MGP,i} \times f_{INP,i}) \quad \text{Equation (4)}$$

Where:

$\Delta SOC_i$  = Loss of soil organic carbon in land stratum  $i$ , (t C)

$A_{SOC,i}$  = Area of land stratum  $i$ ; (ha)

$SOC_{REF,i}$  = Reference SOC stock applicable to land stratum  $i$ , (t C/ha)

<sup>4</sup> Based on 2006 IPCC Guidelines for National Greenhouse Gas Inventories. For details, see appendix 4.3.



|             |   |   |
|-------------|---|---|
| $f_{LUB,i}$ | = | Relative stock change factor for land-use in the baseline in stratum $i$  |
| $f_{MGB,i}$ | = | Relative stock change factor for land management in the in stratum $i$  |
| $f_{INB,i}$ | = | Relative stock change factor for input in the baseline in stratum $i$   |
| $f_{LUP,i}$ | = | Relative stock change factor for land-use in the project in stratum $i$   |
| $f_{MGP,i}$ | = | Relative stock change factor for land management in the project in stratum $i$                                  |
| $f_{INP,i}$ | = | Relative stock change factor for input in the project in stratum $i$  |
| $i$         | = | Strata of areas of land   |
| 1.21        | = | Conservativeness factor accounting for the uncertainties in the values in Tables 2-4 in appendix 1 <sup>5</sup> |

17. The values of relative stock change factors shall be determined according to Tables 2-4 in appendix 1 of this tool.<sup>6</sup>

18. After the first crediting period of the project, the value of  $PE_{SOC,y}$  shall be 0.

### 5.1.2. Emissions resulting from soil fertilization and management

19. Emissions resulting from soil fertilisation and management are estimated as follows:

$$PE_{SF,y} = q_{N,y} \times A_{FTM,y} \times EF_{FT} \quad \text{Equation (5)}$$

Where:

|             |   |  |
|-------------|---|--|
| $PE_{SF,y}$ | = | Emissions resulting from soil fertilization and management, in year $y$ (t CO <sub>2</sub> e)  |
| $q_{N,y}$   | = | Rate of nitrogen applied, in year $y$ (t N/ha)   |
| $A_{FTM,y}$ | = | Area of land subjected to soil fertilization and management, in year $y$ (ha)  |
| $EF_{FT}$   | = | Aggregate emission factor for N <sub>2</sub> O and CO <sub>2</sub> emissions resulting from production and application of nitrogen (t CO <sub>2</sub> e/(t N)). A default value of 13.3 t CO <sub>2</sub> e/(t N) <sup>7</sup> shall be used |

### 5.1.3. Emissions resulting from soil amendment

20. Emissions resulting from soil amendment (liming) are estimated as follows:

<sup>5</sup> According to FCCC/SBSTA/2003/10/Add.2/6. For details, see appendix 1.

<sup>6</sup> Project proponents are encouraged to suggest revisions for this tool with alternative procedures (e.g. monitoring) to determine the relative stock change.

<sup>7</sup> Based on 2006 IPCC Guidelines for National Greenhouse Gas Inventories. For details, see appendix 1.

$$PE_{SA,y} = q_{LM,y} \times A_{LM,y} \times EF_{LM} + q_{DL,y} \times A_{DL,y} \times EF_{DL} \quad \text{Equation (6)}$$

Where:

|             |  |
|-------------|--|
| $PE_{SA,y}$ | = Emissions resulting from soil amendment by liming, in year y (t CO <sub>2</sub> e)   |
| $q_{LM,y}$  | = Rate of application of limestone, in year y (t/ha)   |
| $A_{LM,y}$  | = Area of land in which limestone is applied, in year y (ha)   |
| $EF_{LM}$   | = Emission factor for CO <sub>2</sub> emissions from limestone application (t CO <sub>2</sub> e/(t limestone)). A default value of 0.12 t CO <sub>2</sub> e/(t limestone) <sup>8</sup> shall be used |
| $q_{DL,y}$  | = Rate of application of dolomite, in year y (t/ha)  |
| $A_{DL,y}$  | = Area of land in which dolomite is applied, in year y (ha)  |
| $EF_{DL}$   | = Emission factor for CO <sub>2</sub> emissions from dolomite application (t CO <sub>2</sub> e (t dolomite)). A default value of 0.13 t CO <sub>2</sub> e/(t dolomite) <sup>9</sup> shall be used    |

## 5.2. Emissions resulting from energy consumption

21. Emissions resulting from energy consumption are estimated by following the provisions in the methodological tools “Tool to calculate project or leakage CO<sub>2</sub> emissions from fossil fuel combustion” and “Tool to calculate baseline, project and/or leakage emissions from electricity consumption”.

## 5.3. Emissions resulting from clearance or burning of biomass

22. Emissions resulting from clearance or burning of biomass are estimated as follows:

$$PE_{BB,y} = \frac{44}{12} \times 0.47 \times \sum_i A_{FR,i,z} \times b_i \times (1.07 + R_i) \quad \text{Equation (7)}$$

Where:

|                 |  |
|-----------------|--|
| $PE_{BB,y}$     | = Emissions resulting from burning of biomass, in year y (t CO <sub>2</sub> e)   |
| $\frac{44}{12}$ | = Factor for converting units from t C to t CO <sub>2</sub> e; dimensionless   |
| 0.47            | = Default value of carbon fraction of biomass burnt; <sup>10</sup> dimensionless   |
| 1.07            | = Factor to account for non-CO <sub>2</sub> emissions from biomass burning; <sup>11</sup> dimensionless. If biomass is cleared without using open fire, then this factor is set equal to 1 (one) |

<sup>8</sup> 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Vol 4, Ch 11, Eq 11.12.

<sup>9</sup> Ibid.

<sup>10</sup> 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Vol 4, Ch 4 Table 4.3.

|              |   |  |
|--------------|---|--|
| $A_{FR,i,y}$ | = | Area of stratum $i$ of land subjected to fire in year $y$ (ha)   |
| $b_i$        | = | Fuel biomass consumption per hectare in stratum $i$ of land subjected to fire (t dry matter/ha)  |
| $R_i$        | = | Root-shoot ratio (i.e. ratio of below-ground biomass to above-ground biomass) for stratum $i$ of land subjected to fire; dimensionless |
| $i$          | = | Strata of areas of land  |

## 6. Leakage due to diversion of biomass residues

### 6.1. Determination of the alternative scenario of the biomass residues in absence of the project activity

23. For the alternative fate of biomass residues that will be used in the underlying CDM project activity, the alternative scenarios in absence of the project activity shall include, inter alia:

- (a) B1: The biomass residues are dumped or left to decay mainly under aerobic conditions. This applies, for example, to dumping and decay of biomass residues on fields;
- (b) B2: The biomass residues are dumped or left to decay under clearly anaerobic conditions. This applies, for example, to landfills which are deeper than five meters. This does not apply to biomass residues that are stock-piled or left to decay on fields;
- (c) B3: The biomass residues are burnt in an uncontrolled manner without utilizing it for energy purposes;
- (d) B4: The biomass residues are used for electricity generation in power-only plant configuration at the project site in new and/or existing power plants;
- (e) B5: The biomass residues are used for power and/or heat generation in existing or new power plants at other sites;
- (f) B6: The biomass residues are used for other energy purposes, such as the generation of bio-fuels;
- (g) B7: The biomass residues are used for non-energy purposes, e.g. as fertilizer or as feedstock in processes (e.g. in the pulp and paper industry);
- (h) B8: The primary source of the biomass residues and/or their fate in the absence of the project activity cannot be clearly identified.<sup>12</sup>

<sup>11</sup> Based on 2006 IPCC Guidelines for National Greenhouse Gas Inventories. For details, see appendix 1.

<sup>12</sup> For example, this scenario can be used if biomass residues are purchased from a market, or biomass residues retailers, or if processed biomass is purchased from biomass processing plants which are not included in the project boundary.

24. Project proponents may choose to combine some or all relevant biomass types when determining the fate of biomass residues, and treat the combined types as one, for instance in the biomass availability determination. This combinations shall be documented transparently in the CDM-PDD and remain consistent throughout the crediting period.
25. When defining plausible and credible alternative scenarios for the fate of biomass residues, the guidance below shall be followed:
- (a) If the biomass residues processing (drying, pelletization, shredding, briquetting, etc.) is not included in the project boundary, the processed biomass obtained from that plant should be considered as B8 above;
  - (b) The alternative scenario for the biomass residues should be separately identified for different categories of biomass residues, covering the whole amount of biomass residues supposed to be used in the project activity along the crediting period;
  - (c) A category of biomass residues is defined by three attributes: (1) its type (i.e. bagasse, rice husks, empty fruit bunches, etc.); (2) its source (e.g. produced on-site, obtained from an identified biomass residues producer, obtained from a biomass residues market, etc.); and (3) its alternative scenario in the absence of the project activity (Scenarios B above);
  - (d) Explain and document transparently in the CDM-PDD, using a table similar to Table 1 in Appendix 2, what quantities of which biomass residues categories are used in which installation(s) under the project activity and what is their alternative scenario;
  - (e) For biomass residues categories for which scenarios B1, B2 or B3 are deemed a plausible alternative scenario, project participants shall demonstrate that this is a realistic and credible alternative scenario. Towards this end one of the following procedures should be applied for the combined amount of biomass identified:
    - (i) Demonstrate that there is an abundant surplus of the biomass residue in the region of the project activity which is not utilized. For this purpose, demonstrate that the quantity of that type of biomass residues annually available in the region is at least 25 per cent larger than the quantity of biomass residues which is utilized annually in the region (e.g. for energy generation or as feedstock), including the project facility;
    - (ii) Demonstrate for the sites from where biomass residues are sourced that the biomass residues have not been collected or utilized (e.g. as fuel, fertilizer or feedstock) but have been dumped and left to decay, land-filled, left in the field to decay after harvest,<sup>13</sup> or burnt without energy generation (e.g. field burning). This approach is only applicable to biomass residues categories for which project participants can clearly identify the site from where the biomass residues are sourced;

---

<sup>13</sup> Project proponents shall demonstrate the fraction of biomass which exceeds the function of refertilising the soil, as only this part of the biomass may be considered unutilised.

- (iii) In case abundance of biomass in the region cannot be demonstrated, the alternative use of the biomass shall be considered unknown (B8) and result in leakage emissions.

26. If during the crediting period, new categories of biomass residues of the type B1, B2 or B3 are used in the project activity which were not listed at the validation stage, for example due to new sources of biomass residues, the alternative scenario for those types of biomass residues should be assessed using the procedures outlined in this tool for each new category of biomass residues.

## 6.2. Calculation of Leakage due to diversion of biomass residues

27. The main potential source of leakage due to biomass residues is an increase in emissions from fossil fuel combustion or other sources due to diversion of biomass residues from other uses to the project plant as a result of the project activity. The alternative scenarios for biomass residues for which this potential leakage is relevant are B4, B5, B6, B7 and B8.

28. The actual leakage emissions in each of these cases may differ significantly and depend on the specific situation of each project activity. For that reason, a simplified approach is used in this tool: it is assumed that an equivalent amount of fossil fuels, on energy basis, would be used if biomass residues are diverted from other users, no matter what the use of biomass residues would be in the alternative scenario.

29. Therefore, for the categories of biomass residues whose alternative scenario has been identified as B4, B5, B6, B7 or B8, project participants shall calculate leakage emissions as follows:

$$LE_{BR,y} = EF_{CO_2,LE} \times \sum_n BR_{PJ,n,y} \times NCV_{n,y} \quad \text{Equation (8)}$$

Where:

|                |   |  |
|----------------|---|--|
| $LE_{BR,y}$    | = | Leakage emissions in year $y$ (t CO <sub>2</sub> e)  |
| $EF_{CO_2,LE}$ | = | CO <sub>2</sub> emission factor of the most carbon intensive fossil fuel used in the country (t CO <sub>2</sub> /GJ)         |
| $BR_{PJ,n,y}$  | = | Quantity of biomass residues used in the project site and included in the project boundary in year $y$ (tonnes on dry-basis) |
| $NCV_{n,y}$    | = | Net calorific value of the biomass residues of category $n$ in year $y$ (GJ/tonne of dry matter)                             |
| $n$            | = | Categories of biomass residues for which B4, B5, B6, B7 or B8 has been identified as the alternative scenario                |

30. The determination of  $BR_{PJ,n,y}$  shall be based on the monitored amounts of biomass residues used in facilities included in the project boundary.

## 7. Leakage due to shift of pre-project activities

31. Project proponents are advised to avoid pre-project activities from being shifted outside the project boundary, to avoid indirect land use changes as a result of the project activity. Rather, project proponents are encouraged to include in the project boundary the land in which the pre-project activities will take place after the project implementation.
32. No leakage due to shift of pre-project activities occurs if one of the following two conditions apply:
- (a) The plantation area was abandoned land prior to the implementation of the project activity;
  - (b) The plantation area was used prior to the implementation of the project area but the pre-project land use of the plantation area will be accommodated for, providing at least the same level of service during the project activity, within the land area included in the project boundary. The project area may be expanded to accommodate for this condition. This could be achieved, inter alia, in the following ways:
    - (i) At least the same number of cattle as prior to the implementation of the project activity will continue being grazed during the project activity within the land area included in project boundary;
    - (ii) Due to more efficient farming practice, the pre-project crops can be grown on a smaller area, which is included in the land area included in the project boundary, to achieve the same level of annual production of crops, freeing land for the dedicated plantation;
    - (iii) Settlements are not removed from the land area included in the project boundary.
33. Project participants should assess the possibility of leakage from the displacement of activities or people considering the following indicators:
- (a) Percentage of families/households of the community involved in or affected by the project activity displaced (from within to outside of the project boundary) due to the project activity;
  - (b) Percentage of total production of the main product (e.g. meat, corn) within the project boundary displaced due to the generation of renewable biomass.
34. If the value of both indicators is lower than 10 per cent, then leakage from this source is assumed to be zero.
35. For project activities which fall below the small-scale threshold, if the value of any of these two indicators is higher than 10 per cent and less than or equal to 50 per cent, then leakage shall be equal to 15 per cent of the difference between baseline emissions and project emissions.
36. If the value of either of these two indicators is larger than 50 per cent for project activities which fall below the small-scale threshold, or 10 per cent otherwise, then this tool is not applicable and a new procedure must be submitted for the approval of the Board.

## 8. Monitoring methodology

### 8.1. Data and parameters not monitored

37. Data and parameters not monitored have been provided along with the relevant equation in the preceding section of the tool. Furthermore, for projects including biomass cultivation, the following parameter shall be known:

**Data / Parameter table 1.**

|                                  |  |
|----------------------------------|--|
| <b>Data / Parameter:</b>         | <b>Pre-project land use</b>  |
| Data unit:                       | variable   |
| Description:                     | Service level of the pre-project land use  |
| Source of data:                  | Land management records, records of the relevant local authority, stakeholders interviews etc. |
| Measurement procedures (if any): | -  |
| Any comment:                     | -  |

**Data / Parameter table 2.**

|                                  |  |
|----------------------------------|--|
| <b>Data / Parameter:</b>         | <b>Biomass residues categories and quantities used in the project activity</b>   |
| Data unit:                       | (a) Type (i.e. bagasse, rice husks, empty fruit bunches, etc.);<br>(b) Source (e.g. produced on-site, obtained from an identified biomass residues producer, obtained from a biomass residues market, etc.);<br>(c) Fate in the absence of the project activity (Scenario B);<br>(d) Use in the project scenario   |
| Description:                     | Explain and document transparently in the CDM-PDD, using a table similar to table 1 in appendix 2 which quantities of which biomass residues categories are used in which installation(s) under the project activity and what is their alternative scenario.<br>The last column of table 1 in appendix 2 corresponds to the quantity of each category of biomass residues (tonnes on dry-basis). These quantities should be updated every year of the crediting period so as to reflect the actual use of biomass residues in the project scenario. These updated values should be used for leakage calculations, if the determined alternative fate indicates associated leakage emissions. Along the crediting period, new categories of biomass residues (i.e. new types, new sources, with different fate) can be used in the project activity. In this case, a new line should be added to the table. If those new categories are of the type B1, B2 or B3, the alternative scenario for those types of biomass residues should be assessed using the procedures outlined in the guidance provided in the procedure for the determination of the alternative scenario |
| Measurement procedures (if any): | -  |
| Any comment:                     | -  |

## 8.2. Data and parameters monitored

38. Data and parameters relevant to projects including biomass cultivation are monitored as described in the following data description tables:

**Data / Parameter table 3.**

|   |   |
|---|---|
| <b>Data / Parameter:</b>                | $A_{SOC,i}$   |
| <b>Data unit:</b>                       | ha  |
| <b>Description:</b>                     | Area of land stratum <i>i</i>   |
| <b>Source of data:</b>                  | Measurement by project participants   |
| <b>Measurement procedures (if any):</b> | Standard land area measurement methods applicable in the host party                             |
| <b>Monitoring frequency:</b>            | Annual  |
| <b>QA/QC procedures:</b>                | Check that standard land area measurement methods applicable in the host party country are used |
| <b>Any comment:</b>                     | -   |

**Data / Parameter table 4.**

|   |   |
|---|---|
| <b>Data / Parameter:</b>                | $q_{N,y}$   |
| <b>Data unit:</b>                       | t N/ha  |
| <b>Description:</b>                     | Rate of nitrogen applied, in year <i>y</i>  |
| <b>Source of data:</b>                  | Land management records maintained by project participants and fertiliser composition information from supplier, study or independent laboratory. Alternatively, the default conservative value of 0.20 t N/ha per year may be used |
| <b>Measurement procedures (if any):</b> | -   |
| <b>Monitoring frequency:</b>            | Annual  |
| <b>QA/QC procedures:</b>                | Cross-check records of applied quantities with purchase receipts and inventory  |
| <b>Any comment:</b>                     | Nitrogen applied through the following methods shall be added up to arrive at this value: (i) synthetic fertilisers; (ii) organic manure; (iii) return of the residues or cover crops   |

**Data / Parameter table 5.**

|   |   |
|---|---|
| <b>Data / Parameter:</b>                | $A_{FTM,y}$   |
| <b>Data unit:</b>                       | ha  |
| <b>Description:</b>                     | Area of land subjected to soil fertilization and management, in year <i>y</i> |
| <b>Source of data:</b>                  | Measurement by project participants   |
| <b>Measurement procedures (if any):</b> | Standard land area measurement methods applicable in the host party           |
| <b>Monitoring frequency:</b>            | Annual  |



|                   |  |
|-------------------|--|
| QA/QC procedures: | Check that standard land area measurement methods applicable in the host party are used  |
| Any comment:      | Areas receiving one or more of the following inputs shall be added up to arrive at this value: (i) synthetic fertilisers; (ii) organic manure; (iii) return of the residues or cover crops |

**Data / Parameter table 6.**

|                                  |  |
|----------------------------------|--|
| Data / Parameter:                | $q_{L,M,y}$  |
| Data unit:                       | t/ha   |
| Description:                     | Rate of application of limestone, in year $y$                                  |
| Source of data:                  | Land management records maintained by project participants                     |
| Measurement procedures (if any): | -  |
| Monitoring frequency:            | Annual   |
| QA/QC procedures:                | Cross-check records of applied quantities with purchase receipts and inventory |
| Any comment:                     | -  |

**Data / Parameter table 7.**

|                                  |   |
|----------------------------------|---|
| Data / Parameter:                | $A_{L,M,y}$   |
| Data unit:                       | ha  |
| Description:                     | Area of land in which limestone is applied, in year $y$                                 |
| Source of data:                  | Measurement by project participants   |
| Measurement procedures (if any): | Standard land area measurement methods applicable in the host party                     |
| Monitoring frequency:            | Annual  |
| QA/QC procedures:                | Check that standard land area measurement methods applicable in the host party are used |
| Any comment:                     | -   |

**Data / Parameter table 8.**

|                                  |  |
|----------------------------------|--|
| Data / Parameter:                | $q_{DL,y}$   |
| Data unit:                       | t/ha   |
| Description:                     | Rate of application of dolomite, in year $y$                                   |
| Source of data:                  | Land management records maintained by project participants                     |
| Measurement procedures (if any): | -  |
| Monitoring frequency:            | Annual   |
| QA/QC procedures:                | Cross-check records of applied quantities with purchase receipts and inventory |
| Any comment:                     | -  |

**Data / Parameter table 9.**

|   |   |
|---|---|
| <b>Data / Parameter:</b>                | $A_{DL,y}$  |
| <b>Data unit:</b>                       | ha  |
| <b>Description:</b>                     | Area of land in which dolomite is applied, in year $y$                                  |
| <b>Source of data:</b>                  | Measurement by project participants   |
| <b>Measurement procedures (if any):</b> | Standard land area measurement methods applicable in the host party                     |
| <b>Monitoring frequency:</b>            | Annual  |
| <b>QA/QC procedures:</b>                | Check that standard land area measurement methods applicable in the host party are used |
| <b>Any comment:</b>                     | -   |

**Data / Parameter table 10.**

|   |   |
|---|---|
| <b>Data / Parameter:</b>                | $Q_{FF,i,y}$  |
| <b>Data unit:</b>                       | l   |
| <b>Description:</b>                     | Quantity of fossil fuel of type $i$ consumed in year $y$  |
| <b>Source of data:</b>                  | Land management records maintained by project participants. Alternatively, the default conservative value of 50 litre of diesel equivalent per hectare per year may be used   |
| <b>Measurement procedures (if any):</b> | -   |
| <b>Monitoring frequency:</b>            | Annual  |
| <b>QA/QC procedures:</b>                | -   |
| <b>Any comment:</b>                     | Fossil fuels used in the following activities shall be added up to arrive at this value: (i) use of farm machinery e.g. tractors, harvesters; (ii) treatment, pumping, and application of water for irrigation; (iii) use of machinery in transport and application of inorganic fertilisers, organic fertilisers, soil amending materials (e.g. limestone, dolomite) |

**Data / Parameter table 11.**

|   |   |
|---|---|
| <b>Data / Parameter:</b>                | $Q_{EL,y}$  |
| <b>Data unit:</b>                       | MWh   |
| <b>Description:</b>                     | Quantity of electricity consumed in year $y$  |
| <b>Source of data:</b>                  | Land management records maintained by project participants or dedicated electricity meters  |
| <b>Measurement procedures (if any):</b> | -   |
| <b>Monitoring frequency:</b>            | Annual  |
| <b>QA/QC procedures:</b>                | -   |
| <b>Any comment:</b>                     | Electricity used in the following activities shall be added up to arrive at this value: (i) use of farm machinery run on electricity; (ii) treatment, pumping, and application of water for irrigation; (iii) transport and application of inorganic fertilisers, organic fertilisers, soil amending materials (e.g. limestone, dolomite) |

**Data / Parameter table 12.**

|   |   |
|---|---|
| <b>Data / Parameter:</b>                | $A_{FR,i,y}$  |
| <b>Data unit:</b>                       | ha  |
| <b>Description:</b>                     | Area of stratum <i>i</i> of land subjected to fire in year <i>y</i>                     |
| <b>Source of data:</b>                  | Measurement by project participants   |
| <b>Measurement procedures (if any):</b> | Standard land area measurement methods applicable in the host party                     |
| <b>Monitoring frequency:</b>            | Annual  |
| <b>QA/QC procedures:</b>                | Check that standard land area measurement methods applicable in the host party are used |
| <b>Any comment:</b>                     | -   |

**Data / Parameter table 13.**

|   |   |
|---|---|
| <b>Data / Parameter:</b>                | $b_i$   |
| <b>Data unit:</b>                       | t dry matter/ha   |
| <b>Description:</b>                     | Fuel biomass consumption per hectare in stratum <i>i</i> of land subjected to fire  |
| <b>Source of data:</b>                  | Measurement by project participants. Alternatively, the default 'average above-ground biomass content in forest' values from Table 3A.1.4 of the Good Practice Guidance for Land Use, Land-Use Change and Forestry (IPCC-GPG-LULUCF 2003) |
| <b>Measurement procedures (if any):</b> | Measurement may be carried out through sample plots   |
| <b>Monitoring frequency:</b>            | Annual  |
| <b>QA/QC procedures:</b>                | If sample plots are used, the estimated mean value should not have an uncertainty of greater than 10 per cent at 90 per cent confidence level   |
| <b>Any comment:</b>                     | -   |

**Data / Parameter table 14.**

|   |  |
|---|--|
| <b>Data / Parameter:</b>                | $R_i$  |
| <b>Data unit:</b>                       | Dimensionless  |
| <b>Description:</b>                     | Root-shoot ratio (i.e. ratio of below-ground biomass to above-ground biomass) for stratum <i>i</i> of land subjected to fire                               |
| <b>Source of data:</b>                  | Measurement by project participants. Alternatively, the default values from Table 4.4 of the 2006 IPCC Guidelines for National GHG Inventories may be used |
| <b>Measurement procedures (if any):</b> | Measurement may be carried out through sample plots  |
| <b>Monitoring frequency:</b>            | Annual   |
| <b>QA/QC procedures:</b>                | If sample plots are used, the estimated mean value should not have an uncertainty of greater than 10% at 90% confidence level                              |
| <b>Any comment:</b>                     | -  |

**Data / Parameter table 15.**

|                                  |  |
|----------------------------------|--|
| <b>Data / Parameter:</b>         | <b>Land use</b>  |
| Data unit:                       | variable   |
| Description:                     | Service level of the project land use  |
| Source of data:                  | -  |
| Measurement procedures (if any): | -  |
| Monitoring frequency:            | Annually   |
| QA/QC procedures:                | -  |
| Any comment:                     | The service level of the project land use shall at least provide the pre-project service level |

39. For projects utilising biomass residues, in the absence of relevant data/parameter tables in the methodology, the following tables shall be used.

**Data / Parameter table 16.**

|                                  |  |
|----------------------------------|--|
| <b>Data / Parameter:</b>         | <b>BR<sub>P,J,n,y</sub></b>  |
| Data unit:                       | tonnes on dry-basis  |
| Description:                     | Quantity of biomass residues of category <i>n</i> used in facilities which are located at the project site and included in the project boundary in year <i>y</i>   |
| Source of data:                  | On-site measurements   |
| Measurement procedures (if any): | Use weight meters. Adjust for the moisture content in order to determine the quantity of dry biomass   |
| Monitoring frequency:            | Data monitored continuously and aggregated as appropriate, to calculate emissions reductions   |
| QA/QC procedures:                | Cross-check the measurements with an annual energy balance that is based on purchased quantities and stock changes   |
| Any comment:                     | The biomass residue quantities used should be monitored separately for (a) each type of biomass residue (e.g.) and each source (e.g. produced on-site, obtained from biomass residues suppliers, obtained from a biomass residues market, obtained from an identified biomass residues producer, etc.) |

**Data / Parameter table 17.**

|                                  |   |
|----------------------------------|---|
| <b>Data / Parameter:</b>         | <b>NCV<sub>n,y</sub></b>  |
| Data unit:                       | GJ/tonnes on dry-basis  |
| Description:                     | Net calorific value of biomass residues of category <i>n</i> in year <i>y</i>   |
| Source of data:                  | On-site measurements  |
| Measurement procedures (if any): | Measurements shall be carried out at reputed laboratories and according to relevant international standards. Measure the NCV on dry-basis |
| Monitoring frequency:            | At least every six months, taking at least three samples for each measurement   |

|                   |  |
|-------------------|--|
| QA/QC procedures: | Check the consistency of the measurements by comparing the measurement results with measurements from previous years, relevant data sources (e.g. values in the literature, values used in the national GHG inventory) and default values by the IPCC. If the measurement results differ significantly from previous measurements or other relevant data sources, conduct additional measurements. Ensure that the NCV is determined on the basis of dry biomass |
| Any comment:      | -  |

**Data / Parameter table 18.**

|                                  |  |
|----------------------------------|--|
| Data / Parameter:                | EF <sub>CO<sub>2</sub>,LE</sub>  |
| Data unit:                       | t CO <sub>2</sub> /GJ  |
| Description:                     | CO <sub>2</sub> emission factor of the most carbon intensive fuel used in the country  |
| Source of data:                  | Identify the most carbon intensive fuel type from the national communication, other literature sources (e.g. IEA). Possibly consult with the national agency responsible for the national communication/GHG inventory. If available, use national default values for the CO <sub>2</sub> emission factor. Otherwise, IPCC default values may be used |
| Measurement procedures (if any): | -  |
| Monitoring frequency:            | Annually   |
| QA/QC procedures:                | -  |
| Any comment:                     | -  |

## Appendix 1. Default values for biomass cultivation

**Table 1. Default reference SOC stocks (SOCREF) for mineral soils (tC/ha in 0–30 cm depth)**

| Climate region        | HAC soils <sup>14</sup> | LAC soils <sup>15</sup> | Sandy soils <sup>16</sup> | Spodic soils <sup>17</sup> | Volcanic soils <sup>18</sup> |
|-----------------------|-------------------------|-------------------------|---------------------------|----------------------------|------------------------------|
| Boreal                | 68                      | NA                      | 10                        | 117                        | 20                           |
| Cold temperate, dry   | 50                      | 33                      | 34                        | NA                         | 20                           |
| Cold temperate, moist | 95                      | 85                      | 71                        | 115                        | 130                          |
| Warm temperate, dry   | 38                      | 24                      | 19                        | NA                         | 70                           |
| Warm temperate, moist | 88                      | 63                      | 34                        | NA                         | 80                           |
| Tropical, dry         | 38                      | 35                      | 31                        | NA                         | 50                           |
| Tropical, moist       | 65                      | 47                      | 39                        | NA                         | 70                           |
| Tropical, wet         | 44                      | 60                      | 66                        | NA                         | 130                          |
| Tropical montane      | 88                      | 63                      | 34                        | NA                         | 80                           |

DRAFT

<sup>14</sup> Soils with high activity clay (HAC) minerals are lightly to moderately weathered soils, which are dominated by 2:1 silicate clay minerals (in the World Reference Base for Soil Resources (WRB) classification these include Leptosols, Vertisols, Kastanozems, Chernozems, Phaeozems, Luvisols, Alisols, Albeluvisols, Solonetz, Calcisols, Gypsisols, Umbrisols, Cambisols, Regosols; in USDA classification includes Mollisols, Vertisols, high-base status Alfisols, Aridisols, Inceptisols).

<sup>15</sup> Soils with low activity clay (LAC) minerals are highly weathered soils, dominated by 1:1 clay minerals and amorphous iron and aluminium oxides (in WRB classification includes Acrisols, Lixisols, Nitisols, Ferralsols, Durisols; in USDA classification includes Ultisols, Oxisols, acidic Alfisols).

<sup>16</sup> Includes all soils (regardless of taxonomic classification) having >70 per cent sand and <8 per cent clay, based on standard textural analyses (in WRB classification includes Arenosols; in USDA classification includes Psamments).

<sup>17</sup> Soils exhibiting strong podzolization (in WRB classification includes Podzols; in USDA classification Spodosols).

<sup>18</sup> Soils derived from volcanic ash with allophanic mineralogy (in WRB classification Andosols; in USDA classification Andisols).

**Table 2. Relative stock change factors for different management activities on cropland<sup>19</sup>**

| Factor type             | Level   | Temperature regime            | Moisture regime   | Factor value | Description and criteria   |
|-------------------------|---|-------------------------------|-------------------|--------------|--|
| Land use ( $f_{LU}$ )   | Long-term cultivated                                      | Temperate/Boreal              | Dry               | 0.80         | Area has been continuously managed for crops for more than 20 years  |
|                         |   |                               | Moist             | 0.69         |  |
|                         |   | Tropical                      | Dry               | 0.58         |  |
|                         |   |                               | Moist/Wet         | 0.48         |  |
| Tropical montane        | n/a   | 0.64                          |                   |              |  |
| Land use ( $f_{LU}$ )   | Short-term cultivated (<20 years) or set aside (<5 years) | Temperate/Boreal and Tropical | Dry               | 0.93         | Area has been managed for crops for less than 20 years and/or the area is cropland that has been in a fallow state for less than five years at any point during the last 20 years                |
|                         |   |                               | Moist/Wet         | 0.82         |  |
|                         |   | Tropical montane              | n/a               | 0.88         |  |
| Management ( $f_{MG}$ ) | Full tillage  | All                           | Dry and Moist/Wet | 1.00         | Substantial soil disturbance with full inversion and/or frequent (within year) tillage operations. At planting time, little (e.g. <30 per cent) of the surface is covered by residues            |
| Management ( $f_{MG}$ ) | Reduced tillage   | Temperate/Boreal              | Dry               | 1.02         | Primary and/or secondary tillage but with reduced soil disturbance (usually shallow and without full soil inversion). Normally leaves surface with >30 per cent coverage by residues at planting |
|                         |   |                               | Moist             | 1.08         |  |
|                         |   | Tropical                      | Dry               | 1.09         |  |
|                         |   |                               | Moist/Wet         | 1.15         |  |
|                         |   | Tropical montane              | n/a               | 1.09         |  |
| Management ( $f_{MG}$ ) | No-tillage  | Temperate/Boreal              | Dry               | 1.10         | Direct seeding without primary tillage, with only minimal soil disturbance in the seeding zone. Herbicides are typically used for weed control   |
|                         |   |                               | Moist             | 1.15         |  |
|                         |   | Tropical                      | Dry               | 1.17         |  |
|                         |   |                               | Moist/Wet         | 1.22         |  |
|                         |   | Tropical montane              | n/a               | 1.16         |  |

<sup>19</sup> Adapted from 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4, Table 5.5.

**Table 3. Relative stock change factors for different levels of nutrient input on cropland<sup>20</sup>**

| Factor type        | Level               | Temperature regime            | Moisture regime   | Factor value | Description and criteria  |
|--------------------|---------------------|-------------------------------|-------------------|--------------|---|
| Input ( $f_{IN}$ ) | Low                 | Temperate/Boreal              | Dry               | 0.95         | There is removal of residues (via collection or burning), or frequent bare-fallowing, or production of crops yielding low residues (e.g. vegetables, tobacco, cotton), or no mineral fertilization or N-fixing crops  |
|                    |                     |                               | Moist             | 0.92         |   |
|                    |                     | Tropical                      | Dry               | 0.95         |   |
|                    |                     |                               | Moist/Wet         | 0.92         |   |
| Tropical montane   | n/a                 | 0.94                          |                   |              |   |
| Input ( $f_{IN}$ ) | Medium              | All                           | Dry and Moist/Wet | 1.00         | All crop residues are returned to the field. If residues are removed then supplemental organic matter (e.g. manure) is added. Additionally, mineral fertilization or N-fixing crop rotation is practised  |
| Input ( $f_{IN}$ ) | High without manure | Temperate/Boreal and Tropical | Dry               | 1.04         | Represents significantly greater crop residue inputs over medium C input cropping systems due to additional practices, such as production of high residue yielding crops, use of green manures, cover crops, improved vegetated fallows, irrigation, frequent use of perennial grasses in annual crop rotations, but without manure applied |
|                    |                     |                               | Moist/Wet         | 1.11         |   |
|                    |                     | Tropical Montane              | n/a               | 1.08         |   |
|                    |                     |                               |                   |              |   |
| Input ( $f_{IN}$ ) | High with manure    | Temperate/Boreal and Tropical | Dry               | 1.37         | Represents significantly higher C input over medium C input cropping systems due to an additional practice of regular addition of animal manure   |
|                    |                     | -                             | Moist/Wet         | 1.44         |   |
|                    |                     | Tropical Montane              | n/a               | 1.41         |   |

<sup>20</sup> Ibid.



**Table 4. Relative stock change factors ( $f_{LU}$ ,  $f_{MG}$ , and  $f_{IN}$ ) for grassland management<sup>21</sup>**

| Factor type  | Level                         | Climate regime   | Factor value | Description  |
|--|-------------------------------|------------------|--------------|--|
| Land use ( $f_{LU}$ )                                      | All                           | All              | 1.00         | All permanent grassland is assigned a land-use factor of 1   |
| Management ( $f_{MG}$ )                                    | Non-degraded grassland        | All              | 1.00         | Non-degraded and sustainably managed grassland, but without significant management improvements  |
| Management ( $f_{MG}$ )                                    | Moderately degraded grassland | Temperate/Boreal | 0.95         | Overgrazed or moderately degraded grassland, with somewhat reduced productivity (relative to the native or nominally managed grassland) and receiving no management inputs       |
|  |                               | Tropical         | 0.97         |  |
|  |                               | Tropical Montane | 0.96         |  |
| Management ( $f_{MG}$ )                                    | Severely degraded             | All              | 0.70         | Implies major long-term loss of productivity and vegetation cover, due to severe mechanical damage to the vegetation and/or severe soil erosion                                  |
| Management ( $f_{MG}$ )                                    | Improved grasslands           | Temperate/Boreal | 1.14         | Represents grassland which is sustainably managed with moderate grazing pressure and that receive at least one improvement (e.g. fertilization, species improvement, irrigation) |
|  |                               | Tropical         | 1.17         |  |
|  |                               | Tropical Montane | 1.16         |  |
| Input ( $f_{IN}$ )<br>(applied only to improved grassland) | Medium                        | All              | 1.00         | All grassland without input of fertilizers is assigned an input factor of 1  |
|  | High                          | All              | 1.11         | Grasslands with direct application of fertilizers (organic or inorganic) beyond what is required to be classified as improved grassland  |

<sup>21</sup> Adapted from 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4, Table 6.2.

## Appendix 2. Example identification of alternative uses

1. Consider a project activity which includes the installation of a new biomass-only power plant, and the retrofit of an existing co-fired biomass-fossil-fuel power plant, which has historically used rice husks, produced on-site. Suppose that the project activity will use two types of biomass residues, rice husks (historical use plus an additional amount) and diverse agricultural residues (as additional biomass residues compared to the historical situation). Further consider that the rice husks used in the project would come from two different sources, on-site production and off-site supply from an identified rice mill. Presumably, the rice husks produced on-site would have been partly used on-site for electricity generation and partly be dumped in the determined alternative scenario. The rice husks procured off-site would have been dumped in the determined alternative scenario. The diverse agricultural residues are purchased from a biomass retailer. For this example, four categories of biomass residues should be considered in the subsequent analysis, as illustrated in Table 1.
2. The last column of Table 1 corresponds to the quantity of each category of biomass residues (tonnes). For the determination of the alternative scenario, at the validation stage, an ex ante estimation of these quantities should be provided. These quantities should be updated every year of the crediting period as part of the monitoring plan so as to reflect the actual use of biomass residues in the project scenario. These updated values should be used for emissions reductions calculations. Along the crediting period, new categories of biomass residues (i.e. new types, new sources, with different fate) can be used in the project activity. In this case, a new line should be added to the table.

**Table 1. Table for biomass residues categories**

| Biomass residues category ( <i>k</i> ) | Biomass residues type | Biomass residues source                   | Biomass residues fate in the absence of the project activity | Biomass residues use in project scenario             | Biomass residues quantity (tonnes) |
|--|-----------------------|---|--|--|------------------------------------|
| 1                                      | Rice husks            | On-site production                        | Electricity generation on-site (B4)                          | Electricity generation on-site (biomass-only boiler) | See comments above                 |
| 2                                      | Rice husks            | On-site production                        | Dumped (B1)  | Electricity generation on-site (biomass-only boiler) | See comments above                 |
| 3                                      | Rice husks            | Off-site from an identified rice mill     | Dumped (B1)  | Electricity generation on-site (biomass-only boiler) | See comments above                 |
| 4                                      | Agricultural residues | Off-site from a biomass residues retailer | Unidentified (B8)  | Electricity generation on-site (co-fired boiler)     | See comments above                 |

## Appendix 3. Explanation of factors used

1. In equation (3), the factor to account for soil N<sub>2</sub>O emissions associated with loss of soil organic carbon is calculated following volume 4 chapters 3 and 11 of 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Loss of SOC leads to associated mineralisation of N in the soil, leading to N<sub>2</sub>O emissions:
  - (a) The mineralised *N* can be calculated using equation (11.1) and (11.10), with  $EF_1=0.01$ ,  $EF_5 = 0.0075$  and  $Frac_{LEACH} = 0.3$ , results in total 0.01225 tN-N<sub>2</sub>O/tN;
  - (b) Using equation (11.8) of the IPCC guidelines, in which *R* is set to 10 tSOC/tN, results in 0.001225 tN-N<sub>2</sub>O/tSOC;
  - (c) Converting to t CO<sub>2</sub>e/tSOC by multiplying with 298 (GWP<sub>N<sub>2</sub>O</sub>) and dividing by 28/44 (Weight of *N* in N<sub>2</sub>O) results in 0.574 t CO<sub>2</sub>e/tSOC;
  - (d) Dividing by 44/12 (mass ratio of CO<sub>2</sub> and C) to convert to t CO<sub>2</sub>e/t CO<sub>2</sub> (dimensionless factor) results in 0.156 t CO<sub>2</sub>e released in N<sub>2</sub>O for each t CO<sub>2</sub> released from SOC.
2. In equation (4), the factor to account for the IPCC default factor was derived from evaluating worse-case scenario, i.e. worse uncertainties, in the used factors:
  - (a) Reviewing the IPCC data,  $SOC_{REF}$  has error estimate of 90 per cent (IPCC table 2.3, table note), whereas the various *f* factors have error estimate of up to 50 per cent (IPCC tables 5.4 and 6.2). These are two sigma estimates, equivalent to 95 per cent confidence interval;
  - (b) Converting them to 90 per cent confidence interval (equivalent to 1.282 sigma), which is deemed appropriate for the tool, by multiplying with 1.282/2, results in  $SOC_{REF}$  uncertainty of 58 per cent, and the various *f* factors in uncertainties of 32 per cent;
  - (c) Adding the root-mean-square of these error estimates result in total 70 per cent error (Note the *f* uncertainties have each half the weight of the  $SOC_{REF}$  error estimate, due to the addition in the equation);
  - (d) As  $SOC_{REF}$  always has error estimate of 58 per cent, total error estimate has range of 58–70 per cent;
  - (e) The error estimates being in the uncertainty band of 50-100 per cent, result in a corrective factor of 1.21 according to FCCC/SBSTA/2003/10/Add.2/6.

- 
3. In equation (5), the default value of the aggregate emission factor for N<sub>2</sub>O and CO<sub>2</sub> emissions resulting from production and application of nitrogen, is calculated following volume 4 chapters 3 and 11 of 2006 IPCC Guidelines for National Greenhouse Gas Inventories, as well as based on Wood and Cowie (2004) and Swaminathan (2004). The calculation is for ammonium nitrate, which is considered conservative:<sup>22</sup>
- (a) Direct and indirect N<sub>2</sub>O emissions (emissions associated to the fertiliser application on the soil) calculated, using IPCC equations (11.1), (11.9) and (11.10), with  $EF_1 = 0.01$ ,  $EF_4 = 0.01$ ,  $EF_5 = 0.0075$ ,  $Frac_{GASF} = 0.1$  and  $Frac_{LEACH} = 0.3$ , results in total 0.01325 tN-N<sub>2</sub>O/tN. This is converted, by multiplying with 298 (GWP<sub>N<sub>2</sub>O</sub>) dividing by 28/44 (Weight of N in N<sub>2</sub>O), to 6.20 t CO<sub>2</sub>e/tN;
  - (b) Emissions from synthetic fertiliser production, including fuel, feedstocks and emissions during production, calculated based on Wood and Cowie (2004) and Swaminathan (2004), taken for ammonium nitrate, a conservative fertiliser, is 7.1 t CO<sub>2</sub>e/tN;
  - (c) Adding the above emissions results in 13.3 t CO<sub>2</sub>e/tN.
4. In equation (78), the factor to account for non-CO<sub>2</sub> emissions from biomass burning was calculated using the values in table 2.5, volume 4 chapter 2 of 2006 IPCC Guidelines for National Greenhouse Gas Inventories:
- (a) Taking the values for savannah and grassland, which are most conservative;
  - (b)  $(1613 \text{ g CO}_2 + 2.3 \text{ g CH}_4 \times 25 (\text{GWP}_{\text{CH}_4}) + 0.21 \text{ g N}_2\text{O} \times 298 (\text{GWP}_{\text{N}_2\text{O}})) / 1613 \text{ g CO}_2 = 1.07$ .

---

<sup>22</sup> As a comparison, the calculation is repeated for organic fertiliser:

- (a) Direct and indirect N<sub>2</sub>O emissions calculated similarly to ammonium nitrate, but with  $Frac_{GASM} = 0.2$  resulting in 6.67 t CO<sub>2</sub>e/tN;
- (b) No emissions from fertiliser production are considered, resulting in total 6.67 t CO<sub>2</sub>e/tN.

As an additional comparison, the calculation is also repeated for synthetic urea:

- (a) Direct and indirect N<sub>2</sub>O identical to ammonium nitrate equals 6.20 t CO<sub>2</sub>e/tN;
- (b) Emissions from urea production, from the same source as ammonium nitrate, is 1.70 t CO<sub>2</sub>e/tN;
- (c) Emissions from urea applications (carbon released from the urea decomposition) calculated as 0.429 tC/tN, which is the C/N mass ratio in urea. This is converted, by dividing by 12/44 (Weight of C in CO<sub>2</sub>), to 1.57 t CO<sub>2</sub>e/tN;
- (d) Adding the above emissions results in 9.5 t CO<sub>2</sub>e/tN.

### Document information

---

| <i>Version</i> | <i>Date</i>     | <i>Description</i>   |
|----------------|-----------------|--|
| Draft 02.0     | 3 November 2014 | MP 65, annex 10<br>A call for public input will be launched on the draft revised tool.<br>Revision to simplify and streamline the requirements for accounting for leakage emissions from use of biomass residues/biomass from cultivation. |
| 01.0           | 4 October 2013  | EB 75, Annex 11<br>Initial adoption.   |

---

Decision Class: Regulatory  
Document Type: Tool  
Business Function: Methodology  
Keywords: biomass, leakage

---

DRAFT